

Salt, I. P. and Hardie, G. (2017) AMP-activated protein kinase: an ubiquitous signalling pathway with key roles in the cardiovascular system. *Circulation Research*, 120(11), pp. 1825-1841.
(doi: [10.1161/CIRCRESAHA.117.309633](https://doi.org/10.1161/CIRCRESAHA.117.309633))

This is the author's final accepted version.

There may be differences between this version and the published version.
You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/139812/>

Deposited on: 13 April 2017

AMP-ACTIVATED PROTEIN KINASE – A UBIQUITOUS SIGNALLING
PATHWAY WITH KEY ROLES IN THE CARDIOVASCULAR SYSTEM

Ian P. Salt¹ and D. Grahame Hardie²

¹Institute of Cardiovascular & Medical Sciences, College of Medical, Veterinary & Life Sciences,
Davidson Building, University of Glasgow, Glasgow, G12 8QQ, Scotland, UK

²Division of Cell Signalling & Immunology, School of Life Sciences, University of Dundee,
Dundee, DD1 5EH, Scotland, UK

Short title: AMPK: roles in cardiovascular system

KEYWORDS: AMPK, AMP-activated protein kinase, heart, heart defects (congenital),
metabolism, vasculature

ABSTRACT

The AMP-activated protein kinase (AMPK) is a key regulator of cellular and whole body energy homeostasis, which acts to restore energy homoeostasis whenever cellular energy charge is depleted. Over the last two decades, it has become apparent that AMPK regulates a number of other cellular functions and has specific roles in cardiovascular tissues, acting to regulate cardiac metabolism and contractile function as well as promoting anti-contractile, anti-inflammatory and anti-atherogenic actions in blood vessels. In this review, we will discuss the role of AMPK in the cardiovascular system, including the molecular basis of mutations in AMPK that alter cardiac physiology and the proposed mechanisms by which AMPK regulates vascular function under physiological and pathophysiological conditions.

The AMP-activated protein kinase (AMPK) is the central component of a signalling pathway that regulates the switch between anabolism and catabolism, as well as many other aspects of cell function¹⁻³. In this review we focus on the physiological roles of AMPK within the cardiovascular system, although we will start by discussing general features that apply in all mammalian cell types. Readers are also referred to two other recent reviews of the role of AMPK in cardiovascular disease^{4, 5}.

STRUCTURE AND REGULATION OF AMPK

Occurrence of subunit isoforms

AMPK exists universally as heterotrimeric complexes containing catalytic α subunits and regulatory β and γ subunits, which occur as multiple isoforms ($\alpha 1/\alpha 2$; $\beta 1/\beta 2$; $\gamma 1/\gamma 2/\gamma 3$) encoded by distinct genes. These could give rise in principle to twelve heterotrimeric combinations, although specific combinations appear to be favoured in specific cell types. For example, although skeletal muscle expresses mRNAs encoding all seven subunit isoforms, assays of immunoprecipitated isoforms suggest that AMPK activity in human skeletal muscle is accounted for by just three combinations: $\alpha 1\beta 2\gamma 1$, $\alpha 2\beta 2\gamma 1$ and $\alpha 2\beta 2\gamma 3$ ⁶.

The domain organization of the seven AMPK subunit isoforms are shown in Fig. 1, and a representation of a crystal structure for the human $\alpha 1\beta 2\gamma 1$ heterotrimer⁷ is shown in Fig. 2. Similar structures of the $\alpha 2\beta 1\gamma 1$ ⁸ and $\alpha 1\beta 1\gamma 1$ ⁹ complexes are available.

The α subunits

The α subunits contain a kinase domain (α -KD) at their N-termini, with the small N-lobe (yellow in Figs.1 and 2) and larger C-lobe (green) typical of all protein kinases. The substrate Mg.ATP²⁻ binds in a deep cleft between these lobes, occupied in the structure in Fig. 2 by the non-specific kinase inhibitor staurosporine. Isolated α -KDs are only significantly active after phosphorylation by upstream kinases at a conserved threonine, usually referred to as Thr172. Thr172 phosphorylation is a good marker for AMPK activity, and Western blotting using phosphospecific antibodies against this site is often used as a semi-quantitative measure of AMPK activity. However, this ignores the effects of allosteric activation of AMPK, a problem that can be overcome by also addressing the phosphorylation state of a downstream target such as acetyl-CoA carboxylase (ACC).

The α -KD is followed by a small globular domain termed the autoinhibitory domain (AID, orange) that, when AMP is not bound to the γ subunit, binds to both the N- and C-lobes of the KD and thus causes inhibition⁷. The AID is connected to the C-terminal domain (α -CTD, red) by the α -linker (blue), a region of extended polypeptide that interacts with the γ subunit when AMP is bound at site 3 (as is the case in Fig. 2), thus pulling the AID away from this inhibitory position.

The β subunits

Two well-conserved regions within the β subunits are the C-terminal domain (β -CTD, silver-grey), which forms the “core” of the heterotrimeric complex, and the central carbohydrate-binding module (β -CBM, silver-blue). The β -CBM is related to carbohydrate-binding domains usually occurring in enzymes that metabolize starch or glycogen, which localize those enzymes on their polysaccharide substrate. Consistent with this, the β -CBM causes a proportion of AMPK in cells to bind to the surface of glycogen particles¹⁰⁻¹². In all three structures of mammalian heterotrimers, the β -CBM interacts with the N-lobe of the α subunit KD, with its glycogen-binding site (defined by binding of the oligosaccharide β -cyclodextrin) at the top in the view of Fig. 2.

What is the function of glycogen binding by AMPK? The muscle and liver isoforms of glycogen synthase, which are also bound to glycogen particles, are physiological targets for AMPK^{13, 14}, and one function may be to co-localize AMPK with them. The β -CBM is also of interest because the cleft between it and the N-lobe of the α subunit forms the binding site for activators such as A769662 and 991⁸. This site has been termed the Allosteric Drug and Metabolite (ADaM) site¹⁵, and is discussed further below.

The γ subunits

All γ subunits contain four tandem repeats of a sequence motif known as a CBS motif. In other proteins, pairs of CBS motifs often form binding sites for regulatory ligands containing adenosine¹⁶; in the AMPK- γ subunits they bind the regulatory nucleotides AMP, ADP or and ATP¹⁷. Two views of a structure for the γ 1 subunit, containing three molecules of bound AMP¹⁸, are shown in Fig. 3. The two pairs of repeats (CBS1:CBS2 and CBS3:CBS4) assemble in a pseudosymmetrical “head-to-head” manner. The γ subunit thus forms a structure like a flattened disc (seen from different faces in Fig. 3) with one CBS repeat in each quadrant, generating four potential ligand-binding clefts in the centre. However, one of these appears to be unused, perhaps because conserved

aspartate residues in CBS1, CBS3 and CBS4 that bind the ribose ring of adenine nucleotides in sites 1, 3 and 4 are absent from CBS2¹⁸. Interestingly, mutation of any of these three aspartate residues interferes with multiple effects of AMP on kinase activity¹⁹, suggesting that occupancy of all three sites may be required for full activation.

Identity of upstream kinases

Identifying the upstream kinases that phosphorylate Thr172 was a difficult challenge, eventually solved by genome-wide biochemical screens that identified three upstream kinases in budding yeast^{20, 21}. The mammalian kinases with catalytic domains most closely related to these were LKB1 and the Ca²⁺/calmodulin-dependent kinase CaMKK2 (CaMKK β), and evidence was soon obtained that both could act as physiological upstream kinases in mammalian cells²²⁻²⁷. The discovery that LKB1 was an upstream kinase for AMPK was particularly interesting, because LKB1 had previously been identified to be a tumor suppressor²⁸. Phosphorylation of Thr172 by CaMKK2²⁵⁻²⁷ represents a mechanism by which hormones that increase cytosolic Ca²⁺ can activate AMPK in the absence of energy stress.

GENERAL FEATURES OF AMPK REGULATION

Regulation by the canonical energy stress mechanism

It might be expected that a system that monitors cellular energy status would sense ATP and ADP but, interestingly, all metabolic enzymes known to directly monitor cellular energy charge (glycogen phosphorylase, 6-phosphofructo-1-kinase, fructose-1,6-bisphosphatase) primarily sense AMP and ATP, as does AMPK. The principal source of AMP in cells is thought to be the adenylate kinase reaction ($2\text{ADP} \leftrightarrow \text{ATP} + \text{AMP}$), which appears to operate close to equilibrium in most cells so that the AMP:ATP ratio will vary as the square of the ADP:ATP ratio²⁹. The former is therefore a more sensitive indicator of falling energy status than the latter.

Binding of AMP activates AMPK by three complementary mechanisms, of which the first two are mimicked by ADP at higher concentration, while all three are antagonized by ATP: (i) inhibition of Thr172 dephosphorylation by protein phosphatases; (ii) promotion of Thr172 phosphorylation by LKB1; and (iii) allosteric activation^{30, 31}. The structural model shown in Fig. 2 suggests a mechanism, for which there is now supporting evidence⁷, to explain mechanisms (i) and

(iii). When AMP is bound at site 3, the α -linker interacts with the surface of the γ subunit containing that site (Fig. 2)^{7, 32}. AMPK heterotrimers contain two rather distinct regions, the “catalytic module” (β -CBM, KD, AID, top/front section in Fig. 2) and the “nucleotide-binding module” (γ subunit, β -CTD, α -CTD, bottom/rear section in Fig. 2). The “hinge” connecting them is the α -linker, and release of the α -linker on binding of ATP rather than AMP at site 3 is envisaged to allow the two modules to move apart, causing the AID to rotate back into its inhibitory position behind the α -KD. This conformational change would also increase accessibility of Thr172 to protein phosphatases, which in the AMP-bound conformation of Fig. 2 is around the back, located in a deep cleft between the two modules. This model therefore explains not only how AMP binding at site 3 causes allosteric activation, but also why it protects against Thr172 dephosphorylation, with binding of ATP antagonizing both effects.

Multiple mechanisms of pharmacological activation of AMPK

A selection of compounds commonly used to activate AMPK experimentally are listed in Table 1. Those in Class 1, including the antidiabetic drug metformin and berberine (derived from traditional Chinese medicine) inhibit Complex I of the mitochondrial respiratory chain, while those in class 2, including 2-deoxyglucose, inhibit glycolysis. Both classes activate AMPK indirectly by increasing cellular AMP:ATP ratios³³. These compounds are frequently used to activate AMPK experimentally, and studies utilizing them are described later in this review. However, because they work by depleting cellular ATP, they should not be regarded as specific AMPK activators and any results obtained with them should ideally be followed up using genetic approaches, such as the use of AMPK knockouts.

The third class of activator includes the widely used compound 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR), a nucleoside that is taken up into cells and converted to the equivalent nucleotide, ZMP. An important caveat here is that although ZMP is an AMP analog that mimics the effects of AMP to activate AMPK, it is about 50-fold less potent than AMP itself³⁴. Because ZMP accumulates to millimolar concentrations inside cells AICAR does cause AMPK activation, but ZMP also has off-target effects. For example, it is known to mimic the effects of AMP on phosphorylase³⁵ and fructose-1,6-bisphosphatase³⁶ as well as AMPK. Another problem with AICAR is that it is an adenosine analogue and, while it does not appear to bind directly to

adenosine receptors, in incubated cell systems it competes with endogenous adenosine for reuptake into cells by adenosine transporters, so can have adenosine-like effects³⁷.

A more specific activator that works via a related mechanism (class 4) is C13, which is taken up by cells and converted by cellular esterases to the AMP analog C2, a very potent AMPK activator³⁸ although only for α 1- and not α 2-containing complexes³⁹. Despite its selectivity for the α 1 isoform, C2 binds to the γ subunit. Although the AMP and C2 binding sites overlap, they are not identical⁴⁰.

The other molecules of choice for selective activation of AMPK are those binding at the ADaM site located between the β -CBM and the N-lobe on the α subunit, such as A769662 and 991 (class 5)⁸. All compounds binding this site are more potent activators of complexes containing AMPK- β 1 rather than β 2, and most are essentially β 1-selective (Table 1). Using AMPK phosphorylated on Thr172, these activators cause a modest degree of allosteric activation (up to 4-fold) and, by inhibiting dephosphorylation, also promote net Thr172 phosphorylation^{41,42}. More remarkably, they cause a much larger allosteric activation (up to 65-fold) of AMPK that is not phosphorylated on Thr172, and this effect is synergistic with AMP⁴³. At present, almost all of the activators known to bind at this site are synthetic compounds derived from high-throughput screens, although there is much speculation in the field that there is a naturally occurring ligand, hence the appearance of “metabolite” in its name¹⁵. However, no natural ligands occurring in animal cells that bind the ADaM site have yet been found. Salicylate, the natural plant product from which acetyl salicylic acid (ASA, aspirin) was derived, does activate AMPK by binding this site^{9,44}.

It should also be noted that there are currently no specific pharmacological inhibitors of AMPK; compound C (dorsomorphin) is sometimes claimed to be a specific inhibitor, but this is not the case⁴⁵.

TARGETS AND PATHWAYS DOWNSTREAM OF AMPK

Catabolic effects switched on by AMPK

Once activated by energy stress, AMPK switches on catabolic pathways generating ATP, while switching off anabolic pathways and other processes consuming ATP, thus acting to restore energy homeostasis. This topic has been discussed in more detail in previous reviews¹⁻³. Examples of catabolic processes acutely switched on include cellular glucose uptake mediated by GLUT1 and GLUT4. Activation of GLUT1 may occur via phosphorylation of the thioredoxin interacting protein

TXNIP⁴⁶, while enhanced GLUT4 translocation to the plasma membrane in muscle appears to occur, at least in part, by phosphorylation of TBC1D1, which modulates trafficking of GLUT4-containing vesicles⁴⁷. AMPK activation also enhances GLUT4 expression⁴⁸, in part via phosphorylation of the histone deacetylase HDAC5, promoting binding of 14-3-3 proteins and consequent retention of HDAC5, a transcriptional inhibitor, in the cytoplasm^{49, 50}. AMPK can also cause a short-term activation of glycolysis via phosphorylation of PFKFB2⁵¹ and PFKFB3⁵², isoforms of the bifunctional enzyme that synthesizes and breaks down fructose-2,6-bisphosphate, a key allosteric activator of 6-phosphofructo-1-kinase and hence glycolysis. PFKFB2 is expressed in the heart, while PFKFB3 occurs as an “inducible” form whose expression in monocytes and macrophages is induced by inflammatory mediators such as lipopolysaccharide⁵³. Phosphorylation of PFKFB2 or PFKFB3 at equivalent sites near their C-termini increases the synthesis of fructose-2,6-bisphosphate and hence promotes glycolysis during hypoxia in heart and in activated monocytes/macrophages, respectively. This may enhance survival of these cells during periods of hypoxia or ischemia.

Although AMPK can therefore activate glucose uptake and glycolysis in specific cell types, in the longer term it tends to promote instead the more glucose-sparing and energy-efficient oxidative metabolism. In skeletal muscle^{54, 55} and liver⁵⁶, AMPK activates fatty acid oxidation by inhibiting the ACC1/ACC2 isoforms of acetyl-CoA carboxylase to reduce malonyl-CoA, an inhibitor of the uptake of fatty acids into mitochondria. AMPK promotes the expression of TCA cycle enzymes⁵⁷, as well as mitochondrial biogenesis, which it achieves by increasing the expression/activity of the transcriptional co-activator PGC-1 α , either by direct phosphorylation triggering a positive feedback effect on its own expression⁵⁸, or by enhancing its deacetylation by SIRT1⁵⁹. PGC-1 α acts as a co-activator for several transcription factors involved in mitochondrial biogenesis and oxidative metabolism, including myocyte enhancer factor-2 (MEF2), nuclear respiratory factors-1/-2 (NRF1/2), and PPAR- α and - δ ⁶⁰. Despite this evidence that AMPK promotes oxidative metabolism, in mice with a skeletal/cardiac muscle-specific knockout of both AMPK- β subunits, although the mice displayed evidence of dilated cardiomyopathy the rate of glucose and fatty acid oxidation in hearts perfused under normoxic conditions *ex vivo* was normal⁶¹.

A final catabolic pathway switched on by AMPK is autophagy, brought about via direct phosphorylation of the protein kinase that triggers that process, ULK1⁶². By triggering digestion of

cellular contents, autophagy may be critical in enhancing cell survival during periods of acute nutrient starvation, such as during ischemia in cardiac muscle. However, it has also been suggested that excessive autophagy might also contribute to cell damage during ischemia/reperfusion⁶³.

Anabolic pathways switched off by AMPK

As well as its classical roles in inhibiting fatty acid and sterol synthesis³⁴, AMPK inactivates key enzymes and regulatory proteins involved in triglyceride/phospholipid synthesis⁶⁴, glycogen synthesis^{13, 14}, rRNA synthesis⁶⁵, and protein synthesis. The latter is inhibited both at the elongation step by phosphorylation and activation of elongation factor-2 (EF2) kinase⁶⁶, and at the initiation step by inactivation of the mechanistic target-of-rapamycin complex-1 (mTORC1) by multiple mechanisms^{67, 68}. By inhibiting mTORC1, AMPK would also be expected to oppose hypertrophy in organs such as the heart, which in most cases is considered to be a deleterious process.

ROLE OF AMPK IN THE HEART

Role in the response to cardiac ischemia

It was reported in 1995 that AMPK was activated by no-flow ischemia in perfused rat hearts, an effect associated with high rates of fatty acid oxidation during reperfusion⁶⁹. AMPK is also activated in the heart by increased workload⁷⁰. The role of AMPK in cardiac ischemia has subsequently been addressed using mouse models where AMPK is down-regulated or absent. The first was a transgenic model where an inactive AMPK- $\alpha 2$ subunit was expressed in both skeletal and cardiac muscle. By competing with endogenous α subunits for available β and γ subunits, the over-expressed inactive $\alpha 2$ subunit down-regulates endogenous $\alpha 1$ and $\alpha 2$, and thus acts as a dominant negative mutant. There was no major phenotype under unstressed conditions, but during low-flow ischemia there was a failure to enhance glucose uptake and glycolysis, while during subsequent reperfusion there was less fatty acid oxidation, lower ATP levels and poorer recovery of contractile function⁷¹. The degree of damage to the myocardium was also greater, suggesting that AMPK exerts a cardioprotective effect overall, even though stimulation of fatty acid oxidation during reperfusion may be deleterious⁶⁹. Using the same model, evidence was obtained that AMPK was required for increased autophagy during ischemia⁷².

Next to be studied were conditional knockouts of the upstream kinase LKB1 in both skeletal and cardiac muscle. The basal activity of AMPK- α 2 complexes in the heart was completely abolished, and did not increase in response to no-flow ischemia or anoxia, correlating with a complete lack of ACC phosphorylation at the AMPK site. The AMP:ATP and ADP:ATP ratios were also elevated to a greater extent during no-flow ischemia than in control hearts, confirming that AMPK was protecting the cells against energetic stress. Surprisingly, the activity of AMPK- α 1 complexes was almost unaffected by LKB1 knockout, and still increased in response to ischemia or anoxia⁷³. Broadly similar results were obtained in studies of perfused hearts from mice with a whole body knockout of AMPK- α 2⁷⁴. Interestingly, left ventricular hypertrophy induced by phenylephrine was greatly accentuated in the hearts from AMPK- α 2 knockout mice; this may have been due to less restraint on the mTORC1 pathway, because the basal phosphorylation of p70S6K at the mTORC1 site was elevated, although it did not increase further upon phenylephrine treatment as in the controls⁷⁵.

Mutations in γ 2 and γ 3 subunits causing heart disease and altered glycogen content

The only mutations in any of the seven genes encoding AMPK subunits clearly shown to cause human pathology are those in *PRKAG2*, encoding AMPK- γ 2. These are associated with multiple cardiac disorders, specifically: (i) ventricular pre-excitation; (ii) excessive glycogen storage in cardiomyocytes; (iii) cardiac hypertrophy. Although rare, the *PRKAG2* mutations are autosomal dominant in effect and therefore occur frequently in affected families. They are invariably missense mutations causing amino acid replacements, of which at least 14 have now been reported (Table 2). The affected residues are perfectly conserved between human γ 1, γ 2 and γ 3, and occur in all four CBS repeats. The location of the residues affected in γ 2, and of the three bound molecules of AMP, are mapped in Fig. 3 onto a structure for γ 1¹⁸. Interestingly, six of the mutations (R302Q, H383R, R384T, H530R, R531G and R531Q) affect basic residues whose side chains directly interact with the phosphate groups of one or more molecules of AMP or ATP, while two others also interact either directly (S548) or indirectly (T400) with AMP¹⁸. Other affected residues lie in more peripheral regions of the γ subunit, and it is less obvious why their replacement should affect function. In the context of bacterially expressed CBS repeats from AMPK- γ 2, the R302Q, H383R, T400N, and R531G mutations all reduced the affinity for ATP as well as AMP, with the severity of

the effects increasing in the order R302Q<H383R<T400N<R531G¹⁶. When expressed in mammalian cells as $\alpha 1\beta 1\gamma 2$ heterotrimers, these mutations also reduced allosteric activation by AMP, with the R531G mutation abolishing AMPK activation completely^{16, 76}. The R384T and R531Q mutations were shown later to cause severe effects on AMPK function, similar to or even greater than R531G^{77, 78}. In a comparison of the R531G and R531Q mutations expressed as $\alpha 1\beta 1\gamma 2$ heterotrimers in HEK-293 cells, these mutations not only abolished allosteric activation by AMP, but also caused significant increases in basal Thr172 phosphorylation and activity⁷⁸. This was particularly clear in cells stably expressing the R531G mutant, when clones could be selected in which the level of expression of the WT and mutant was identical; the R531G mutant consistently had a 2-fold higher basal Thr172 phosphorylation and activity although, unlike the wild type, it was completely insensitive to further activation by agents that increase cellular AMP/ADP³³.

Some of the mutations, such as R302Q, can cause relatively mild symptoms so that the patients may not present in the clinic until early adulthood⁷⁹. By contrast, patients with the R531G mutation develop quite severe symptoms during childhood⁸⁰, while the R384T⁷⁷ and R531Q⁷⁸ mutations were only detected *post mortem* in neonates who had died within weeks of birth, and appeared to be *de novo* mutations because neither parent was affected. Consistent with the idea that R531G, R531Q and R384T cause particularly severe forms of the disease, these were the mutations with the largest effects on binding of AMP and ATP to the γ subunit^{16, 77, 78}. It is also interesting that R531 and R384 are involved in the binding of AMP and/or ATP at the critical site, site 3.

What causes the cardiac sequelae of the *PRKAG2* mutations? The reduction of AMP binding to the γ subunit causes reduced activation by AMP and is a loss-of-function effect, whereas the reduction of ATP binding may be responsible for the increased basal activity of the R531G and R531Q mutants and is a gain-of-function effect. Since these mutations are dominant, it seems likely that it is the gain-of-function effect, increased basal activity, that causes most of the pathology; the loss-of-function effect may be compensated for by $\gamma 1$, which is the major isoform expressed in heart, at least in rodents⁸¹. This is an important conclusion because pharmaceutical companies are developing AMPK activators to combat Type 2 diabetes, and it suggests that activation of $\gamma 2$ -containing complexes in the heart might have deleterious effects, as seen with these mutations.

Why does increased basal activity of $\gamma 2$ -containing complexes cause these cardiac phenotypes? AMPK is known to promote glucose uptake in cardiac muscle⁷¹, so increased basal activity would

be expected to cause a high basal uptake, even in the absence of a real demand for glucose. There is good evidence for this scenario from studies of transgenic mice over-expressing the N488I mutation in the heart, which develop ventricular pre-excitation and cardiac hypertrophy similar to the human disorder⁸². By 50 days of age, they have a 20-fold increase in cardiac glycogen compared with controls, accompanied by higher rates of glucose uptake and glycogen synthesis but lower rates of lactate production, suggesting that increased glucose uptake is being directed into glycogen synthesis rather than glycolysis. Glycogen synthase was also much more highly phosphorylated in the transgenic mice (presumably due to the high basal AMPK activity), but cellular glucose-6-phosphate (G6P, an allosteric activator of glycogen synthase that over-rides the effects of phosphorylation) was also elevated >3-fold. Satisfying confirmation of this interpretation came when the N488I mice were crossed with knock-in mice carrying a mutation that renders glycogen synthase insensitive to G6P. The presence of the G6P-insensitive glycogen synthase reversed the high glycogen phenotype of the N488I mice and rescued the ventricular pre-excitation but not the cardiac hypertrophy, suggesting that the former but not the latter is secondary to increased glycogen content⁸³. Supporting the idea that hyper-activation of the mTORC1 pathway was the cause of hypertrophy, mTORC1 targets such as p70S6K and 4EBP1 were more highly phosphorylated in the N488I mice, while treatment with rapamycin reduced, without completely preventing, the hypertrophy. Increased mTORC1 in the N488I mice is actually rather counter-intuitive, because AMPK is normally thought to inhibit that pathway^{67, 68}. However, a recent study has suggested that AMPK can activate mTORC1 under certain circumstances by sustaining the supply of amino acids via autophagy⁸⁴.

Why does abnormally high glycogen content lead to ventricular pre-excitation? During foetal development, the atria and the ventricles become separated by the growth of a fibrous layer called the *annulus fibrosis*, which ensures that the only electrical connection between the two chambers is via the atrioventricular node. In N488I mice the annulus fibrosis appeared to be thin and disrupted in places, and it was suggested that the presence of glycogen-containing vacuoles in myocytes disrupts its formation during fetal development, causing abnormal electrical connections between the atria and ventricles⁸⁵.

Most mouse studies of $\gamma 2$ mutations have involved transgenic mice over-expressing the mutations in human $\gamma 2$. Although these mice do display the key clinical features of the human

syndrome, and mice expressing wild type $\gamma 2$ from the same promoters were used as controls, they are not perfect models because $\gamma 2$ is being over-expressed. Recently, three knock-in mouse models, in which mutations in the mouse gene equivalent to human R302Q, N488I or R531G are expressed globally, were studied^{86, 87}. There was evidence for increased basal AMPK activity in liver and/or muscle of all three strains. The N488I and R531G mice displayed ventricular pre-excitation and modest increases in heart weight, which was associated with large increases in glycogen content in hearts of R531G but not the N488I mice. Both N488I and R531G mice displayed resistance to obesity and hepatic steatosis induced by high-fat diet, but, unlike the N488I mice, the R531G mice also displayed impaired renal function associated with glycogen accumulation, cyst formation, and inflammation and apoptosis in the kidney, particularly when on a high-fat diet⁸⁷. By contrast, in the R302Q mice there was no obvious cardiac phenotype, but the homozygotes developed marked obesity as they aged, with smaller effects in heterozygotes. Obesity appeared to be due mainly to increased food intake through enhanced action of ghrelin, whose effects are mediated by AMPK activation via the CaMKK2 pathway in the hypothalamus^{88, 89}. The homozygotes were also hypoinsulemic, apparently due to reduced glucose-stimulated insulin secretion from the pancreas. Overall, the phenotypes of these mice with knock-in mutations in the *PRKAG2* gene showed surprising variability. Interestingly, heterozygous human carriers of the R302Q mutation, who have a relatively mild cardiac phenotype, also display some evidence of increased obesity, as well as higher fasting glucose and glycated haemoglobin (HbA1c) and lower insulin levels than unaffected siblings⁸⁶. These non-cardiac features of the R302Q mutation only became evident following studies of the mouse model.

Finally, all of the residues mutated in AMPK- $\gamma 2$ are also conserved in $\gamma 1$ and $\gamma 3$ (Table 2). It is interesting that none of them have yet been reported to be mutated in the human $\gamma 1$ gene. However, mutations equivalent to R302Q have been found in both pigs and humans in the $\gamma 3$ isoform, which is predominantly expressed in skeletal muscle⁸¹. In human muscle, $\gamma 3$ appears to be present exclusively as the $\alpha 2\beta 2\gamma 3$ complex, which is the only form of AMPK that is activated during exercise⁶. An R200Q mutation (equivalent to R302Q in human $\gamma 2$) was found to be a relatively common dominantly acting genetic variant in Hampshire pigs, and was associated with a high glycogen content in skeletal muscle; although adversely affecting meat quality, it did not cause obvious clinical problems⁹⁰. Interestingly, in a screen of around 1500 humans, an R225W mutation

(R225 being the human equivalent of R302) was found in two apparently unrelated individuals, once again not associated with any obvious clinical defects; muscles from humans with this mutation had a higher basal AMPK activity, 2-fold higher glycogen content and lower triglyceride content⁹¹. In an *in vitro* study of myotubes from R225W carriers, they had 3-fold higher mitochondrial content and oxidative capacity, and 2-fold higher basal glucose uptake and glycogen synthesis rates than matched controls. The R225W subjects also had a remarkable resistance to fatigue during isometric contractions of the quadriceps⁹², suggesting that this mutation might confer an advantage during endurance exercise because of the high glycogen content.

ROLE OF AMPK IN THE VASCULATURE

In addition to the heart, it has become clear that AMPK has an important role in regulating vascular function, with distinct roles in the functions of vascular endothelial cells (ECs), smooth muscle cells (VSMCs), adventitial cells and vascular immune cells.

Regulation of AMPK in endothelial and vascular smooth muscle cells

AMPK- α 1 accounts for the majority of total AMPK activity in ECs⁹³⁻⁹⁵, yet specific down-regulation of AMPK- α 2 still has marked effects^{96,97}. A wide variety of physiological stimuli have been reported to activate endothelial AMPK, including hypoxia, low glucose and shear stress⁹⁸⁻¹⁰⁰, adiponectin¹⁰¹, angiotensin II and ghrelin^{102,103}. In addition, CaMKK2 mediates AMPK activation by vasoactive molecules including thrombin, VEGF, sphingosine-1-phosphate, bradykinin and estrogen¹⁰⁴⁻¹⁰⁸, suggesting that any stimulus that increases Ca^{2+} in ECs will activate AMPK. Finally, a number of widely used hypoglycemic (metformin, thiazolidinediones, salicylate, DPP4 inhibitors and liraglutide)¹⁰⁹⁻¹¹² and hypocholesterolemic (statins and fenofibrate)^{113,114} drugs activate AMPK in cultured ECs.

Physiological signals that *inhibit* AMPK in ECs include high nutrient concentrations¹¹⁵⁻¹¹⁷, yet the mechanisms by which this inhibition of AMPK in ECs occurs are unclear, although increased PP2A-mediated dephosphorylation of AMPK has been proposed¹¹⁵. Endothelial AMPK may therefore be suppressed by the high levels of nutrients associated with obesity and insulin resistance. As protein kinase C (PKC) activation is associated with over-nutrition, it is interesting that PKC-mediated inhibitory phosphorylation of Ser487 on AMPK- α 1 has been recently reported in ECs¹¹⁸. Conversely, the very first study of AMPK function in ECs demonstrated that AICAR

stimulated fatty acid oxidation¹¹⁹ and AMPK activation was subsequently found to normalise impaired fatty acid oxidation and insulin signaling due to high glucose¹²⁰. Stimulation of fat oxidation may partly underlie the effect of AMPK to antagonize palmitate-mediated endothelial dysfunction, thus protecting against lipotoxicity.

As with ECs, AMPK- α 1 accounts for the majority of AMPK activity in murine and human VSMCs^{121, 122}, and low glucose, adiponectin, estradiol and metformin all activate AMPK¹²³⁻¹²⁶. Angiotensin II has also been reported to acutely activate AMPK¹²⁷, although prolonged activation had no effect¹²⁸. Similar to ECs, high glucose inhibits AMPK in VSMCs, and IGF-1 also inhibits AMPK, most likely via inhibitory phosphorylation of AMPK- α 1 by Akt¹²⁹. Several studies have demonstrated that culture in phosphate/ β -glycerophosphate, used to examine VSMC calcification as described later, also inhibits AMPK^{130, 131}.

AMPK and endothelial NO synthesis

Endothelium-derived NO is a key regulator of vascular function, stimulating VSMC relaxation whilst inhibiting pro-inflammatory signaling, leukocyte adhesion, platelet aggregation, VSMC proliferation and migration associated with pathological vascular remodelling and atherosclerosis¹³². The first indication of a specific vascular role for AMPK came when AMPK was shown to phosphorylate at Ser1177 and activate endothelial NO synthase (eNOS), both in cell-free assays and in ECs^{93, 133}. Multiple mechanisms regulate eNOS activity, including Ca²⁺/calmodulin binding, allosteric and protein:protein interactions, co-factor and substrate availability, phosphorylation, and subcellular localisation¹³². In addition to Ser1177, AMPK phosphorylates eNOS at Ser633, required for AICAR-stimulated NO synthesis in HEK293 cells¹³⁴. NO synthesis requires dimerization of eNOS and sufficient tetrahydrobiopterin (BH₄), in whose absence eNOS generates superoxide instead¹³². Another mechanism by which AMPK improves NO synthesis may be by enhancing BH₄ levels, as AMPK prevents degradation of GTP cyclohydrolase I, which is the rate-limiting enzyme in BH₄ synthesis¹³⁵. AMPK can therefore act by multiple mechanisms to stimulate NO production (Fig. 4), although endothelial AMPK activation is not always associated with eNOS phosphorylation or NO synthesis^{94, 136}. AMPK may also mediate some of the endothelial effects of NO, because NO donors activate AMPK in ECs⁹⁶.

Endothelial AMPK and reactive oxygen species

Inappropriate levels of reactive oxygen species (ROS), in particular superoxide anions generated in response to high concentrations of glucose, lipids and proinflammatory cytokines by NAD(P)H oxidase (Nox), uncoupled eNOS or mitochondrial respiratory chain complexes, have been implicated in vascular disease^{97, 135, 137, 138}. Superoxide reacts with NO to form peroxynitrite, reducing NO bioavailability¹³². AMPK activation in ECs has been widely demonstrated to inhibit ROS formation, increase antioxidant defences and promote mitochondrial biogenesis. Down-regulation of AMPK in ECs increased activity and expression of Nox⁹⁷, whereas AMPK-dependent inhibition of Nox1/2 translocation to the plasma membrane has also been reported¹⁰⁹. The mechanism by which AMPK inhibits Nox remains unclear, but may be secondary to reduced PKC-mediated Nox activation, or NFκB-mediated Nox transcription^{97, 109}. AMPK-dependent inhibition of mitochondrial ROS formation has also been reported in ECs maintained in high glucose¹³⁷, whereas AMPK-mediated stimulation of BH₄ synthesis, as described above, prevents uncoupling of eNOS and superoxide formation¹³⁵. Several groups have reported that AMPK-mediated inhibition of ROS in ECs is associated with increased levels of the antioxidant enzymes superoxide dismutase-2 (SOD2), catalase and thioredoxin^{137, 139, 140}. In ECs, AMPK activation reduces ER stress¹⁴¹, which is tightly linked to oxidative stress and inflammation, whereas silencing of AMPK increases markers of ER stress⁹⁵. Taken together, AMPK activation acts via multiple mechanisms to suppress chronic ROS synthesis in ECs, limiting their damaging actions as well as the sequestration of NO (Fig 4).

AMPK and vascular cell inflammation

The development of endothelial dysfunction and cardiovascular diseases is associated with elevated TNFα, IL-1β and IL-6¹⁴². TNFα and IL-1β stimulate activation of NFκB and the Jun N-terminal kinase (JNK) pathway, while IL-6 signals via Janus kinases (JAKs) leading to phosphorylation of signal transducer and activation of transcription (STAT) proteins¹⁴³. The anti-inflammatory actions of AMPK in ECs were first described when AMPK activation was shown to suppress palmitate- or TNFα-stimulated NFκB activity¹¹⁷. Under basal conditions, NFκB is in an inactive form in the cytoplasm due to binding to IκB whereas, after cytokine stimulation, IκB phosphorylation by IκB kinase (IKK) targets it for degradation, allowing NFκB-mediated transcription of proinflammatory

cytokines, adhesion molecules and chemokines^{143, 144}. NO inhibits endothelial NFκB activity¹⁴⁵, suggesting that AMPK-stimulated NO synthesis would inhibit NFκB, although NO donors do not effectively suppress NFκB in ECs with reduced AMPK activity⁹⁶. AMPK-α2 has been reported to phosphorylate IKKβ *in vitro*, inhibiting IκB phosphorylation and NFκB activation, with reduced IL-1β-stimulated IKK phosphorylation observed in ECs lacking AMPK-α2 but not -α1⁹⁶. Alternatively, AMPK-mediated phosphorylation of the transcriptional co-activator p300 has been proposed to block acetylation and DNA binding by the p65 subunit of NFκB¹⁴⁶. The idea that AMPK inhibits NFκB signalling is reinforced by functional studies demonstrating AMPK-dependent inhibition of NFκB-regulated expression of adhesion molecules and MCP-1^{96, 144}. Fewer studies have investigated the effect of AMPK on proinflammatory JNK and IL-6 signaling, although AICAR and metformin reduce JNK activity in ECs¹⁴⁷, and increased JNK phosphorylation has been reported in ECs lacking AMPK-α2⁹⁵. The mechanism of JNK inhibition is uncertain, although AMPK-dependent inhibition of the upstream kinase MKK4 has been reported in other cells¹⁴⁸. Recently, AMPK-dependent inhibition of IL-6-stimulated JAK-STAT signaling has been demonstrated in ECs, potentially via direct inhibitory phosphorylation of JAK1 by AMPK¹¹⁰. AMPK activation therefore appears to rapidly suppress multiple proinflammatory signaling pathways in ECs, by diverse mechanisms.

AMPK activation in VSMCs inhibits TNFα-stimulated NFκB activity and angiotensin II-stimulated STAT1 activity, as well as reducing expression of inducible NOS and cyclooxygenase-2, and secretion of IL-6 and MCP-1^{149, 150}. Thus, AMPK has anti-inflammatory effects in VSMCs as well as ECs.

AMPK and angiogenesis

Hypoxia, VEGF and adiponectin all stimulate AMPK-dependent EC migration, although there are conflicting reports as to whether this is mediated by NO^{94, 98, 101, 105, 108}. Conversely, down-regulation of AMPK attenuates angiogenesis caused by hypoxia, adiponectin, VEGF or statins, in either tube formation or matrigel plug assays^{94, 98, 101, 113}. Mechanistically, increased UCP2 or SOD2 have been reported to increase angiogenesis in AMPK-deficient ECs^{151, 152}, indicating that down-regulation of ROS may be critical. AMPK activation also stimulates VEGF expression, indicating that AMPK positively influences angiogenesis both by increasing VEGF levels and by increasing

VEGF signaling¹⁵³. As angiogenesis would also consume significant amounts of ATP, AMPK activation might also serve a permissive role, ensuring adequate generation of ATP to permit EC migration and proliferation.

AMPK and VSMC contraction

AMPK has direct anti-contraction effects on VSMCs, because AICAR relaxes aortic rings in an NO- and endothelium-independent manner, an effect lost in AMPK- α 1 knockouts¹²¹. Furthermore, AMPK activation has been reported to affect VSMC contractile signalling, including dephosphorylation of myosin light chain (MLC) and/or myosin phosphatase targeting subunit 1 (MYPT1)^{125, 154}. Mechanistically, MLC/MYPT1 dephosphorylation may be a consequence of AMPK-mediated inhibition and phosphorylation of RhoA at Ser188, causing subsequent inhibition of ROCK¹²⁵. A769662 has also been reported to reduce intracellular Ca^{2+} in VSMCs by increasing sarco/endoplasmic Ca^{2+} ATPase (SERCA) activity, associated with increased phosphorylation of phospholamban on Thr17, which disinhibits SERCA and thereby may underlie SERCA activation and vessel relaxation¹⁵⁵.

AMPK and proliferation, differentiation and migration of VSMCs

Unlike ECs, where AMPK up-regulates proliferation and migration thus supporting angiogenesis, in VSMCs AMPK inhibits proliferation in response to angiotensin II, PDGF and FCS, associated with stimulation of p53 Ser15 phosphorylation and reduced Rb phosphorylation^{128, 156}. VSMCs from mice lacking AMPK- α 2 exhibit increased proliferation, an effect mediated by increased degradation of the cyclin-dependent kinase inhibitor p27^{Kip1} triggered by the ubiquitin E3-ligase Skp2¹⁵⁷. VSMCs retain significant plasticity *in vivo* and can exhibit a synthetic, proliferative phenotype rather than the quiescent, contractile phenotype observed during atherogenesis. AMPK not only suppresses VSMC proliferation, but also inhibits migration and maintains a pro-contraction phenotype^{158, 159}. Inhibition of migration and proliferation are likely to be linked, because increased migration of VSMCs lacking AMPK- α 2 is reported to be Skp2-dependent¹⁵⁹. Furthermore, AICAR limits neointima formation after wire injury of rat femoral arteries, which is likely to reflect the anti-proliferative, anti-migratory actions of AMPK on VSMCs¹²⁷.

AMPK and VSMC calcification

As mentioned above, culture in high phosphate/ β -glycerophosphate concentrations is used experimentally to stimulate calcium deposition in VSMCs. *In vivo*, such vascular calcification is frequently associated with ageing, atherosclerosis and diabetes mellitus. AICAR, adiponectin and metformin all inhibit VSMC calcification *in vitro*^{130, 131, 160}. In recent studies of atherosclerosis-prone *ApoE*^{-/-} mice, deletion of AMPK- α 1 but not AMPK- α 2 caused greater calcification of atherogenic plaques, and levels of the osteogenic transcription factor Runx2¹⁶⁰. Furthermore, metformin reduced atherosclerotic calcification and Runx2 expression in the mice, an effect that was absent in *ApoE*^{-/-} mice lacking AMPK- α 1. The authors of that study further demonstrated that VSMC-specific AMPK- α 1 deletion in *ApoE*^{-/-} mice phenocopied the increased calcification and Runx2 expression, whereas macrophage-specific AMPK- α 1 deletion had no effect. The mechanism underlying the AMPK-mediated inhibition of Runx2 levels was further proposed to be mediated by phosphorylation of PIAS1 (protein inhibitor of activated STAT-1) at Ser510, which acts as a SUMO E3-ligase to trigger Runx2 SUMOylation and degradation¹⁶⁰.

ROLE OF AMPK IN THE VASCULATURE IN VIVO

As many of the actions of AMPK in ECs and VSMCs should have beneficial effects on vascular function (Fig 5), considerable efforts have been made to examine whether AMPK influences vascular tone, remodelling and atherogenesis *in vivo*. Under physiological conditions, vascular AMPK has been reported to be activated *in vivo* by exercise¹⁶¹ and estradiol¹²⁵, and is suppressed by high fat and fructose diets in rodents^{162, 163}. Furthermore, exercise stimulates aortic AMPK phosphorylation in both ECs and VSMCs as assessed by immunohistochemistry¹⁶¹.

AMPK and vascular tone

As described above, AMPK can stimulate NO synthesis by ECs and independently inhibit contractile protein function in VSMCs *in vitro*. In intact arterial vessels, AICAR stimulates vasodilation in a diverse range of vascular beds from several species^{121, 164-167}, an effect greatly attenuated in mice lacking AMPK- α 1¹²¹. The mechanism of AICAR-stimulated vasodilation remains uncertain, and has been variously reported to be endothelium- and NO-dependent^{164, 167}, partially NO- and endothelium-dependent^{165, 166} or NO- and endothelium-independent¹²¹. Using endothelium-specific knockouts it has been proposed that AMPK- α 1 is important for endothelium-

dependent hyperpolarisation-mediated relaxation of resistance arteries¹⁶⁸. Interestingly, resistance arteries exhibited endothelium-independent dilation in response to A769662¹⁵⁵, suggesting a VSMC-mediated effect. It is possible that ECs and VSMCs exhibit differential sensitivities to AICAR and A769662, or that some of these may be AMPK-independent effects. Despite studies indicating that AMPK- α 2 has a less important role in regulation of vascular tone^{121, 168}, AMPK- α 2 knockout mice are hypertensive and exhibit increased contractile responses to phenylephrine in aortic rings¹⁶⁹. Furthermore, impaired bradykinin-dependent vasodilation has been described in EC-specific AMPK- α 2 knockout mice, thought to be due to increased bradykinin degradation by angiotensin-converting enzyme activity¹⁷⁰. AICAR rapidly reduced blood pressure in spontaneously hypertensive rats, but was without effect in normotensive controls¹⁶⁵, while prolonged administration of AICAR reduced systolic blood pressure in obese Zucker rats¹⁷¹. These data support a role for AMPK in the regulation of vascular tone in disease models, but cannot exclude systemic actions on the heart, kidney or other tissues. Surprisingly, whether systemic administration of more selective AMPK activators, such as A769662, 991 or C13, alters vascular tone has yet to be reported. More recently it has become clear that perivascular adipose tissue (PVAT), which is removed in most myography protocols, has a paracrine anti-contractile effect on the underlying vessel. AMPK activity in resistance arteries has been shown to alter the influence of PVAT-derived mediators¹⁷² and the anti-contractile action of PVAT is absent in mice lacking AMPK- α 1, perhaps due to reduced adiponectin secretion¹⁷³. AMPK in ECs, VSMCs and adventitial PVAT may therefore all contribute to the maintenance of vascular tone, but the cell type that mediates actions of AMPK on vascular tone may change, depending on the location of the vessel.

AMPK and atherosclerosis

The potential anti-atherogenic actions of AMPK have been investigated in vascular injury models and atherosclerosis-prone hypercholesterolemic mice. AICAR attenuates post-ischæmic leukocyte rolling and adhesion to the endothelium *in vivo*, an effect lost in AMPK- α 1 or AMPK- α 2 knockout mice¹⁷⁴, and systemic administration of AICAR or metformin reduces atherosclerotic lesion size, macrophage accumulation and inflammation¹⁷⁵⁻¹⁷⁷. Similarly, berberine reduced the severity of atherosclerotic lesions in atherosclerosis-prone mice, an effect attenuated in AMPK- α 2 knockouts¹⁷⁸. AMPK α 2-deficient mice also exhibited increased atherosclerosis⁹⁵, although the

global deficiency makes it difficult to assess whether this was due to a direct effect on vascular tissues. With respect to plaque stability and progression, recent studies have yielded exciting results. Metformin reduced plaque calcification, and mice with a SMC-specific knockout of AMPK- α 1 had exacerbated calcification of atherosclerotic plaques in brachiocephalic arteries, phenocopying the increased calcification observed in global AMPK- α 1 but not AMPK- α 2 knockouts¹⁶⁰. As the clinical use of metformin is associated with reduced macrovascular morbidity and mortality independently of glycemia¹⁷⁹, these results may help to define the pathways involved. Using a ligation model for studying injury-induced neointima stability in brachiocephalic arteries, mice with deletion of AMPK- α 1 but not AMPK- α 2 exhibited more occlusive lesions, with lower collagen and higher macrophage content, indicative of plaque instability¹⁸⁰. Similarly, in a high fat diet model, SMC-specific AMPK- α 2 knockouts exhibited features of unstable plaques, including phenotype switching of VSMCs¹⁵⁸. These data therefore indicate that AMPK activation may not only inhibit atherogenesis, but also inhibit the generation of vulnerable, calcified plaques. Repair of damaged endothelium is also considered important to prevent endothelial dysfunction after injury, and EC-specific expression of constitutively active AMPK has also been reported to promote re-endothelialization in a wire injury model, attributable in part to increased mobilization and incorporation of endothelial progenitor cells¹⁸¹.

Immune cells are involved in all stages of atherosclerosis, and AMPK suppresses inflammatory signalling, monocyte-to-macrophage differentiation and foam cell formation^{175, 177, 182}. Indeed, myeloid-specific AMPK- α 1 knockout mice fed an atherogenic diet on a LDL receptor knockout background have recently been shown to exhibit exacerbated atherosclerosis, with increased plaque macrophage content and inflammatory gene expression¹⁸². All of these studies suggest that AMPK activation limits atherosclerosis, although all of the results are from rodent models, rather than humans. One intriguing question is how, despite only contributing a small fraction of total AMPK activity in vascular cells, specific down-regulation of AMPK- α 2 has such marked effects on atheroma development in mice^{95, 158, 174, 178}, similar to the specific effects noted with respect to vascular tone^{169, 170}. It remains to be determined whether there are isoform-specific substrates or specific subcellular localisations that contribute to the vascular function of AMPK- α 2.

AMPK and pulmonary vascular remodelling

Several studies have identified a critical role for AMPK in the pulmonary vascular remodelling and perivascular inflammation that characterizes pulmonary arterial hypertension (PAH). Metformin reduces pulmonary artery VSMC proliferation in an AMPK-dependent manner¹⁸³, and inhibits the vascular remodelling of pulmonary arteries in rodent models of PAH^{184, 185}. Mice with an EC-specific lack of AMPK also exhibit accelerated pulmonary remodeling¹⁸⁵. Whether the beneficial action of metformin in PAH is mediated by AMPK remains to be determined. By contrast, AMPK activation by hypoxia may contribute to the development of PAH by promoting pulmonary VSMC survival¹⁸⁶, while AMPK-mediated inhibition of the voltage-gated K⁺ channel Kv1.5 may underlie the acute detrimental effects of hypoxia on PAH¹⁸⁷.

Conclusions and Perspectives

AMPK exists as heterotrimeric complexes consisting of catalytic α subunits and regulatory β and γ subunits. AMPK complexes sense the energy status of cells by sensing increases in the cellular AMP:ATP and/or ADP:ATP ratios. AMP, ADP and ATP bind at up to three sites on the γ subunits; binding of AMP and/or ADP causes activation by promoting net phosphorylation at Thr172 within the activation loop on the α subunit, while binding of AMP only causes further allosteric activation. Once activated by energy stress, AMPK switches on catabolic pathways that generate ATP, while switching off cell growth and proliferation and other processes that consume ATP.

AMPK appears to exert a protective effect in rodent heart during ischemic episodes. Inherited and/or *de novo* mutations in the *PRKAG2* gene (encoding AMPK- γ 2) in humans cause heart disease of varying severity characterized by ventricular pre-excitation, excessive cardiac glycogen content, and hypertrophy. The mutations cause an increase in basal AMPK activity, leading to increased glucose uptake that accounts for the first two abnormalities. They also cause a failure of γ 2-containing complexes to be further activated by AMP, which might explain the hypertrophy through unrestrained activity of the mTORC1 pathway.

In blood vessels, AMPK inactivation is associated with anti-contractile, anti-inflammatory and anti-atherogenic actions on both the vascular endothelium and smooth muscle. Exciting recent studies link AMPK activation to increased stability and reduced calcification of atherosclerotic plaques as well as highlighting a potential role for AMPK in pulmonary arterial hypertension.

Given the critical role of AMPK in the regulation of nutrient metabolism, therapies that activate AMPK may not only normalise metabolic dysfunction but also reduce the burden of cardiovascular complications in obesity and type 2 diabetes. Recent studies using vascular tissue-specific deletion of AMPK- α isoforms are beginning to elucidate specific roles for AMPK- α isoforms, yet despite the wealth of research demonstrating the functional cardiovascular consequences of AMPK activation, the AMPK substrates involved and underlying mechanisms in several cases remain poorly defined. Emerging technologies such as phosphoproteomics may prove beneficial in understanding such mechanisms after AMPK up- or down-regulation in cardiovascular tissues in different disease settings and disease models.

Acknowledgements

Recent studies in the DGH laboratory have been supported by a Senior Investigator Award from the Wellcome Trust (097726) and a Programme Grant from Cancer Research UK (C37030/A15101). Recent studies in the IPS laboratory have been supported by Project Grants from the British Heart Foundation (PG/12/1/29276 and PG/13/82/30483) and Diabetes UK (BDA09/0003904 and BDA12/0004489).

References

1. Hardie DG, Schaffer BE, Brunet A. AMPK: an energy-sensing pathway with multiple inputs and outputs. *Trends Cell Biol.* 2016;26:190-201
2. Hardie DG, Ashford ML. AMPK: regulating energy balance at the cellular and whole body levels. *Physiology.* 2014;29:99-107
3. Hardie DG, Ross FA, Hawley SA. AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nature Rev. Mol. Cell Biol.* 2012;13:251-262
4. Daskalopoulos EP, Dufeys C, Beauloye C, Bertrand L, Horman S. AMPK in cardiovascular diseases. *EXS.* 2016;107:179-201
5. Daskalopoulos EP, Dufeys C, Bertrand L, Beauloye C, Horman S. AMPK in cardiac fibrosis and repair: Actions beyond metabolic regulation. *J. Mol. Cell. Cardiol.* 2016;91:188-200
6. Birk JB, Wojtaszewski JF. Predominant α -1/ β -2/ γ -3 AMPK activation during exercise in human skeletal muscle. *J. Physiol.* 2006;577:1021-1032
7. Li X, Wang L, Zhou XE, Ke J, de Waal PW, Gu X, Tan MH, Wang D, Wu D, Xu HE, Melcher K. Structural basis of AMPK regulation by adenine nucleotides and glycogen. *Cell Res.* 2015;25:50-66

- 634 8. Xiao B, Sanders MJ, Carmena D, Bright NJ, Haire LF, Underwood E, Patel BR, Heath RB,
635 Walker PA, Hallen S, Giordanetto F, Martin SR, Carling D, Gamblin SJ. Structural basis of
636 AMPK regulation by small molecule activators. *Nature Commun.* 2013;4:3017
- 637 9. Calabrese MF, Rajamohan F, Harris MS, Caspers NL, Magyar R, Withka JM, Wang H,
638 Borzilleri KA, Sahasrabudhe PV, Hoth LR, Geoghegan KF, Han S, Brown J, Subashi TA,
639 Reyes AR, Frisbie RK, Ward J, Miller RA, Landro JA, Londregan AT, Carpino PA, Cabral S,
640 Smith AC, Conn EL, Cameron KO, Qiu X, Kurumbail RG. Structural basis for AMPK
641 activation: natural and synthetic ligands regulate kinase activity from opposite poles by
642 different molecular mechanisms. *Structure.* 2014;22:1161-1172
- 643 10. Hudson ER, Pan DA, James J, Lucocq JM, Hawley SA, Green KA, Baba O, Terashima T,
644 Hardie DG. A novel domain in AMP-activated protein kinase causes glycogen storage
645 bodies similar to those seen in hereditary cardiac arrhythmias. *Curr. Biol.* 2003;13:861-
646 866
- 647 11. Polekhina G, Gupta A, Michell BJ, van Denderen B, Murthy S, Feil SC, Jennings IG,
648 Campbell, D.J., Witters LA, Parker MW, Kemp BE, Stapleton D. AMPK b-Subunit targets
649 metabolic stress-sensing to glycogen. *Current Biol.* 2003;13:867-871
- 650 12. Bendayan M, Londono I, Kemp BE, Hardie GD, Ruderman N, Prentki M. Association of
651 AMP-activated protein kinase subunits with glycogen particles as revealed in situ by
652 immunoelectron microscopy. *J. Histochem. Cytochem.* 2009;57:963-971
- 653 13. Jorgensen SB, Nielsen JN, Birk JB, Olsen GS, Viollet B, Andreelli F, Schjerling P, Vaulont S,
654 Hardie DG, Hansen BF, Richter EA, Wojtaszewski JF. The α 2-5'AMP-activated protein
655 kinase is a site 2 glycogen synthase kinase in skeletal muscle and is responsive to
656 glucose loading. *Diabetes.* 2004;53:3074-3081
- 657 14. Bultot L, Guigas B, Von Wilamowitz-Moellendorff A, Maisin L, Vertommen D, Hussain N,
658 Beullens M, Guinovart JJ, Foretz M, Viollet B, Sakamoto K, Hue L, Rider MH. AMP-
659 activated protein kinase phosphorylates and inactivates liver glycogen synthase.
660 *Biochem. J.* 2012;443:193-203
- 661 15. Langendorf CG, Kemp BE. Choreography of AMPK activation. *Cell Res.* 2015;25:5-6
- 662 16. Scott JW, Hawley SA, Green KA, Anis M, Stewart G, Scullion GA, Norman DG, Hardie DG.
663 CBS domains form energy-sensing modules whose binding of adenosine ligands is
664 disrupted by disease mutations. *J. Clin. Invest.* 2004;113:274-284
- 665 17. Townley R, Shapiro L. Crystal structures of the adenylate sensor from fission yeast AMP-
666 activated protein kinase. *Science.* 2007;315:1726-1729
- 667 18. Xiao B, Heath R, Saiu P, Leiper FC, Leone P, Jing C, Walker PA, Haire L, Eccleston JF, Davis
668 CT, Martin SR, Carling D, Gamblin SJ. Structural basis for AMP binding to mammalian
669 AMP-activated protein kinase. *Nature.* 2007;449:496-500
- 670 19. Oakhill JS, Chen ZP, Scott JW, Steel R, Castelli LA, Ling N, Macaulay SL, Kemp BE. beta-
671 Subunit myristoylation is the gatekeeper for initiating metabolic stress sensing by AMP-
672 activated protein kinase (AMPK). *Proc. Natl. Acad. Sci. USA.* 2010;107:19237-19241

- 673 20. Hong SP, Leiper FC, Woods A, Carling D, Carlson M. Activation of yeast Snf1 and
674 mammalian AMP-activated protein kinase by upstream kinases. *Proc. Natl. Acad. Sci. U S*
675 *A*. 2003;100:8839-8843
- 676 21. Sutherland CM, Hawley SA, McCartney RR, Leech A, Stark MJ, Schmidt MC, Hardie DG.
677 Elm1p is one of three upstream kinases for the *Saccharomyces cerevisiae* SNF1 complex.
678 *Curr. Biol.* 2003;13:1299-1305
- 679 22. Woods A, Johnstone SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlattner U,
680 Wallimann T, Carlson M, Carling D. LKB1 is the upstream kinase in the AMP-activated
681 protein kinase cascade. *Curr. Biol.* 2003;13:2004-2008
- 682 23. Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Makela TP, Alessi DR, Hardie DG.
683 Complexes between the LKB1 tumor suppressor, STRADA/b and MO25a/b are upstream
684 kinases in the AMP-activated protein kinase cascade. *J. Biol.* 2003;2:28
- 685 24. Shaw RJ, Kosmatka M, Bardeesy N, Hurley RL, Witters LA, DePinho RA, Cantley LC. The
686 tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates
687 apoptosis in response to energy stress. *Proc. Natl. Acad. Sci. USA*. 2004;101:3329-3335
- 688 25. Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM, Frenguelli BG, Hardie DG.
689 Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for
690 AMP-activated protein kinase. *Cell Metab.* 2005;2:9-19
- 691 26. Woods A, Dickerson K, Heath R, Hong SP, Momcilovic M, Johnstone SR, Carlson M,
692 Carling D. Ca²⁺/calmodulin-dependent protein kinase kinase-beta acts upstream of
693 AMP-activated protein kinase in mammalian cells. *Cell Metab.* 2005;2:21-33
- 694 27. Hurley RL, Anderson KA, Franzone JM, Kemp BE, Means AR, Witters LA. The
695 Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase
696 kinases. *J. Biol. Chem.* 2005;280:29060-29066
- 697 28. Alessi DR, Sakamoto K, Bayascas JR. Lkb1-dependent signaling pathways. *Annu. Rev.*
698 *Biochem.* 2006;75:137-163
- 699 29. Hardie DG, Hawley SA. AMP-activated protein kinase: the energy charge hypothesis
700 revisited. *BioEssays*. 2001;23:1112-1119
- 701 30. Gowans GJ, Hawley SA, Ross FA, Hardie DG. AMP is a true physiological regulator of
702 AMP-activated protein kinase by both allosteric activation and enhancing net
703 phosphorylation. *Cell Metab.* 2013;18:556-566
- 704 31. Ross FA, Jensen TE, Hardie DG. Differential regulation by AMP and ADP of AMPK
705 complexes containing different gamma subunit isoforms. *Biochem. J.* 2016;473:189-199
- 706 32. Xiao B, Sanders MJ, Underwood E, Heath R, Mayer FV, Carmena D, Jing C, Walker PA,
707 Eccleston JF, Haire LF, Saiu P, Howell SA, Aasland R, Martin SR, Carling D, Gamblin SJ.
708 Structure of mammalian AMPK and its regulation by ADP. *Nature*. 2011;472:230-233
- 709 33. Hawley SA, Ross FA, Chevtzoff C, Green KA, Evans A, Fogarty S, Towler MC, Brown LJ,
710 Ogunbayo OA, Evans AM, Hardie DG. Use of cells expressing gamma subunit variants to
711 identify diverse mechanisms of AMPK activation. *Cell Metab.* 2010;11:554-565

- 712 34. Corton JM, Gillespie JG, Hawley SA, Hardie DG. 5-Aminoimidazole-4-carboxamide
713 ribonucleoside: a specific method for activating AMP-activated protein kinase in intact
714 cells? *Eur. J. Biochem.* 1995;229:558-565
- 715 35. Longnus SL, Wambolt RB, Parsons HL, Brownsey RW, Allard MF. 5-Aminoimidazole-4-
716 carboxamide 1-beta -D-ribofuranoside (AICAR) stimulates myocardial glycogenolysis by
717 allosteric mechanisms. *Am. J. Physiol.* 2003;284:R936-R944
- 718 36. Vincent MF, Marangos PJ, Gruber HE, Van den Berghe G. Inhibition by AICA riboside of
719 gluconeogenesis in isolated rat hepatocytes. *Diabetes.* 1991;40:1259-1266
- 720 37. Gadalla AE, Pearson T, Currie AJ, Dale N, Hawley SA, Sheehan M, Hirst W, Michel AD,
721 Randall A, Hardie DG, Frenguelli BG. AICA riboside both activates AMP-activated protein
722 kinase and competes with adenosine for the nucleoside transporter in the CA1 region of
723 the rat hippocampus. *J. Neurochem.* 2004;88:1272-1282
- 724 38. Gomez-Galeno JE, Dang Q, Nguyen TH, Boyer SH, Grote MP, Sun Z, Chen M, Craigo WA,
725 van Poelje PD, MacKenna DA, Cable EE, Rolzin PA, Finn PD, Chi B, Linemeyer DL, Hecker
726 SJ, Erion MD. A potent and selective AMPK activator that inhibits de novo lipogenesis.
727 *ACS Med. Chem. Lett.* 2010;1:478-482
- 728 39. Hunter RW, Foretz M, Bultot L, Fullerton MD, Deak M, Ross FA, Hawley SA, Shpiro N,
729 Viollet B, Barron D, Kemp BE, Steinberg GR, Hardie DG, Sakamoto K. Mechanism of
730 action of Compound-13: an alpha1-selective small molecule activator of AMPK. *Chem.*
731 *Biol.* 2014;21:866-879
- 732 40. Langendorf CG, Ngoei KR, Scott JW, Ling NX, Issa SM, Gorman MA, Parker MW, Sakamoto
733 K, Oakhill JS, Kemp BE. Structural basis of allosteric and synergistic activation of AMPK
734 by furan-2-phosphonic derivative C2 binding. *Nat. Commun.* 2016;7:10912
- 735 41. Sanders MJ, Ali ZS, Hegarty BD, Heath R, Snowden MA, Carling D. Defining the
736 mechanism of activation of AMP-activated protein kinase by the small molecule A-
737 769662, a member of the thienopyridone family. *J. Biol. Chem.* 2007;282:32539-32548
- 738 42. Goransson O, McBride A, Hawley SA, Ross FA, Shpiro N, Foretz M, Viollet B, Hardie DG,
739 Sakamoto K. Mechanism of action of A-769662, a valuable tool for activation of AMP-
740 activated protein kinase. *J. Biol. Chem.* 2007;282:32549-32560
- 741 43. Scott JW, Ling N, Issa SM, Dite TA, O'Brien MT, Chen ZP, Galic S, Langendorf CG, Steinberg
742 GR, Kemp BE, Oakhill JS. Small molecule drug A-769662 and AMP synergistically activate
743 naive AMPK independent of upstream kinase signaling. *Chem. Biol.* 2014;21:619-627
- 744 44. Hawley SA, Fullerton MD, Ross FA, Schertzer JD, Chevtzoff C, Walker KJ, Pegg MW,
745 Zibrova D, Green KA, Mustard KJ, Kemp BE, Sakamoto K, Steinberg GR, Hardie DG. The
746 ancient drug salicylate directly activates AMP-activated protein kinase. *Science.*
747 2012;336:918-922
- 748 45. Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevernic I, Arthur JS, Alessi
749 DR, Cohen P. The selectivity of protein kinase inhibitors: a further update. *Biochem J.*
750 2007;408:297-315

46. Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, Shen CH, Wen J, Asara J, McGraw TE, Kahn BB, Cantley LC. AMPK-dependent degradation of TXNIP upon energy stress leads to enhanced glucose uptake via GLUT1. *Mol. Cell.* 2013;49:1167-1175
47. Chen Q, Xie B, Zhu S, Rong P, Sheng Y, Ducommun S, Chen L, Quan C, Li M, Sakamoto K, MacKintosh C, Chen S, Wang HY. A Tbc1d1 Ser231Ala-knockin mutation partially impairs AICAR- but not exercise-induced muscle glucose uptake in mice. *Diabetologia.* 2017;60:336-345
48. Zheng D, MacLean PS, Pohnert SC, Knight JB, Olson AL, Winder WW, Dohm GL. Regulation of muscle GLUT-4 transcription by AMP-activated protein kinase. *J. Appl. Physiol.* 2001;91:1073-1083
49. McGee SL, van Denderen BJ, Howlett KF, Mollica J, Schertzer JD, Kemp BE, Hargreaves M. AMP-activated protein kinase regulates GLUT4 transcription by phosphorylating histone deacetylase 5. *Diabetes.* 2008;57:860-867
50. Mihaylova MM, Vasquez DS, Ravnskjaer K, Denechaud PD, Yu RT, Alvarez JG, Downes M, Evans RM, Montminy M, Shaw RJ. Class IIa histone deacetylases are hormone-activated regulators of FOXO and mammalian glucose homeostasis. *Cell.* 2011;145:607-621
51. Marsin AS, Bertrand L, Rider MH, Deprez J, Beauloye C, Vincent MF, Van den Berghe G, Carling D, Hue L. Phosphorylation and activation of heart PFK-2 by AMPK has a role in the stimulation of glycolysis during ischaemia. *Current Biol.* 2000;10:1247-1255
52. Marsin AS, Bouzin C, Bertrand L, Hue L. The stimulation of glycolysis by hypoxia in activated monocytes is mediated by AMP-activated protein kinase and inducible 6-phosphofructo-2-kinase. *J. Biol. Chem.* 2002;277:30778-30783
53. Rider MH, Bertrand L, Vertommen D, Michels PA, Rousseau GG, Hue L. 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase: head-to-head with a bifunctional enzyme that controls glycolysis. *Biochem. J.* 2004;381:561-579
54. O'Neill HM, Lally JS, Galic S, Thomas M, Azizi PD, Fullerton MD, Smith BK, Pulinkunnil T, Chen Z, Samaan MC, Jorgensen SB, Dyck JR, Holloway GP, Hawke TJ, van Denderen BJ, Kemp BE, Steinberg GR. AMPK phosphorylation of ACC2 is required for skeletal muscle fatty acid oxidation and insulin sensitivity in mice. *Diabetologia.* 2014;57:1693-1702
55. Merrill GM, Kurth E, Hardie DG, Winder WW. AICAR decreases malonyl-CoA and increases fatty acid oxidation in skeletal muscle of the rat. *Am. J. Physiol.* 1997;273:E1107-E1112
56. Fullerton MD, Galic S, Marcinko K, Sikkema S, Pulinkunnil T, Chen ZP, O'Neill HM, Ford RJ, Palanivel R, O'Brien M, Hardie DG, Macaulay SL, Schertzer JD, Dyck JR, van Denderen BJ, Kemp BE, Steinberg GR. Single phosphorylation sites in ACC1 and ACC2 regulate lipid homeostasis and the insulin-sensitizing effects of metformin. *Nat. Med.* 2013;19:1649-1654
57. Winder WW, Holmes BF, Rubink DS, Jensen EB, Chen M, Holloszy JO. Activation of AMP-activated protein kinase increases mitochondrial enzymes in skeletal muscle. *J. Appl. Physiol.* 2000;88:2219-2226

- 791 58. Jager S, Handschin C, St-Pierre J, Spiegelman BM. AMP-activated protein kinase (AMPK)
792 action in skeletal muscle via direct phosphorylation of PGC-1{alpha}. *Proc. Natl. Acad.*
793 *Sci. USA.* 2007;104:12017-12022
- 794 59. Canto C, Gerhart-Hines Z, Feige JN, Lagouge M, Noriega L, Milne JC, Elliott PJ, Puigserver
795 P, Auwerx J. AMPK regulates energy expenditure by modulating NAD(+) metabolism and
796 SIRT1 activity. *Nature.* 2009;458:1056-1060
- 797 60. Lin J, Handschin C, Spiegelman BM. Metabolic control through the PGC-1 family of
798 transcription coactivators. *Cell Metab.* 2005;1:361-370
- 799 61. Sung MM, Zordoky BN, Bujak AL, Lally JS, Fung D, Young ME, Horman S, Miller EJ, Light
800 PE, Kemp BE, Steinberg GR, Dyck JR. AMPK deficiency in cardiac muscle results in
801 dilated cardiomyopathy in the absence of changes in energy metabolism. *Cardiovasc. Res.*
802 2015;107:235-245
- 803 62. Egan DF, Shackelford DB, Mihaylova MM, Gelino S, Kohnz RA, Mair W, Vasquez DS, Joshi
804 A, Gwinn DM, Taylor R, Asara JM, Fitzpatrick J, Dillin A, Viollet B, Kundu M, Hansen M,
805 Shaw RJ. Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects
806 energy sensing to mitophagy. *Science.* 2011;331:456-461
- 807 63. Wang S, Wang C, Yan F, Wang T, He Y, Li H, Xia Z, Zhang Z. N-Acetylcysteine attenuates
808 diabetic myocardial ischemia reperfusion injury through inhibiting excessive autophagy.
809 *Mediators Inflamm.* 2017;2017:9257291
- 810 64. Muoio DM, Seefeld K, Witters LA, Coleman RA. AMP-activated kinase reciprocally
811 regulates triacylglycerol synthesis and fatty acid oxidation in liver and muscle: evidence
812 that sn-glycerol- 3-phosphate acyltransferase is a novel target. *Biochem. J.*
813 1999;338:783-791
- 814 65. Hoppe S, Bierhoff H, Cado I, Weber A, Tiebe M, Grummt I, Voit R. AMP-activated protein
815 kinase adapts rRNA synthesis to cellular energy supply. *Proc. Natl. Acad. Sci. USA.*
816 2009;106:17781-17786
- 817 66. Browne GJ, Finn SG, Proud CG. Stimulation of the AMP-activated protein kinase leads to
818 activation of eukaryotic elongation factor 2 kinase and to its phosphorylation at a novel
819 site, serine 398. *J. Biol. Chem.* 2004;279:12220-12231
- 820 67. Inoki K, Zhu T, Guan KL. TSC2 mediates cellular energy response to control cell growth
821 and survival. *Cell.* 2003;115:577-590
- 822 68. Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, Turk BE, Shaw
823 RJ. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol. Cell.*
824 2008;30:214-226
- 825 69. Kudo N, Barr AJ, Barr RL, Desai S, Lopaschuk GD. High rates of fatty acid oxidation
826 during reperfusion of ischemic hearts are associated with a decrease in malonyl-CoA
827 levels due to an increase in 5'-AMP-activated protein kinase inhibition of acetyl-CoA
828 carboxylase. *J. Biol. Chem.* 1995;270:17513-17520

- 829 70. Coven DL, Hu X, Cong L, Bergeron R, Shulman GI, Hardie DG, Young LH. Physiologic role
830 of AMP-activated protein kinase (AMPK) in the heart: graded activation during exercise.
831 *Am. J. Physiol.* 2003;285:E629-E636
- 832 71. Russell RR, 3rd, Li J, Coven DL, Pypaert M, Zechner C, Palmeri M, Giordano FJ, Mu J,
833 Birnbaum MJ, Young LH. AMP-activated protein kinase mediates ischemic glucose
834 uptake and prevents postischemic cardiac dysfunction, apoptosis, and injury. *J. Clin.*
835 *Invest.* 2004;114:495-503
- 836 72. Matsui Y, Takagi H, Qu X, Abdellatif M, Sakoda H, Asano T, Levine B, Sadoshima J. Distinct
837 roles of autophagy in the heart during ischemia and reperfusion: roles of AMP-activated
838 protein kinase and Beclin 1 in mediating autophagy. *Circ. Res.* 2007;100:914-922
- 839 73. Sakamoto K, Zarrinpashneh E, Budas GR, Pouleur AC, Dutta A, Prescott AR,
840 Vanoverschelde JL, Ashworth A, Jovanovic A, Alessi DR, Bertrand L. Deficiency of LKB1
841 in heart prevents ischemia-mediated activation of AMPK α 2 but not AMPK α 1.
842 *Am. J. Physiol. Endocrinol. Metab.* 2006;290:E780-E788
- 843 74. Zarrinpashneh E, Carjaval K, Beauloye C, Ginion A, Mateo P, Pouleur AC, Horman S,
844 Vaulont S, Hoerter J, Viollet B, Hue L, Vanoverschelde JL, Bertrand L. Role of the α 2-
845 isoform of AMP-activated protein kinase in the metabolic response of the heart to no-
846 flow ischemia. *Am. J. Physiol. Heart Circ. Physiol.* 2006;291:H2875-H2883
- 847 75. Zarrinpashneh E, Beauloye C, Ginion A, Pouleur AC, Havaux X, Hue L, Viollet B,
848 Vanoverschelde JL, Bertrand L. AMPK- α 2 counteracts the development of cardiac
849 hypertrophy induced by isoproterenol. *Biochem. Biophys. Res. Commun.* 2008;376:677-
850 681
- 851 76. Daniel T, Carling D. Functional analysis of mutations in the gamma 2 subunit of AMP-
852 activated protein kinase associated with cardiac hypertrophy and Wolff-Parkinson-
853 White syndrome. *J. Biol. Chem.* 2002;277:51017-51024
- 854 77. Akman HO, Sampayo JN, Ross FA, Scott JW, Wilson G, Benson L, Bruno C, Shanske S,
855 Hardie DG, Dimauro S. Fatal infantile cardiac glycogenosis with phosphorylase kinase
856 deficiency and a mutation in the gamma-2 subunit of AMP-activated protein kinase.
857 *Pediatr. Res.* 2007;62:499-504
- 858 78. Burwinkel B, Scott JW, Buhner C, van Landeghem FK, Cox GF, Wilson CJ, Hardie DG,
859 Kilimann MW. Fatal congenital heart glycogenosis caused by a recurrent activating
860 R531Q mutation in the g2 subunit of AMP-activated protein kinase (PRKAG2), not by
861 phosphorylase kinase deficiency. *Am. J. Hum. Genet.* 2005;76:1034-1049
- 862 79. Thevenon J, Laurent G, Ader F, Laforet P, Klug D, Duva Pentiah A, Gouya L, Maurage CA,
863 Kacet S, Eicher JC, Albuissou J, Desnos M, Bieth E, Duboc D, Martin L, Reant P, Picard F,
864 Bonithon-Kopp C, Gautier E, Binquet C, Thauvin-Robinet C, Faivre L, Bouvagnet P,
865 Charron P, Richard P. High prevalence of arrhythmic and myocardial complications in
866 patients with cardiac glycogenosis due to PRKAG2 mutations. *Europace.* 2016

- 867 80. Gollob MH, Seger JJ, Gollob TN, Tapscott T, Gonzales O, Bachinski L, Roberts R. Novel
868 PRKAG2 mutation responsible for the genetic syndrome of ventricular preexcitation and
869 conduction system disease with childhood onset and absence of cardiac hypertrophy.
870 *Circulation*. 2001;104:3030-3033
- 871 81. Cheung PCF, Salt IP, Davies SP, Hardie DG, Carling D. Characterization of AMP-activated
872 protein kinase α subunit isoforms and their role in AMP binding. *Biochem. J.*
873 2000;346:659-669
- 874 82. Luptak I, Shen M, He H, Hirshman MF, Musi N, Goodyear LJ, Yan J, Wakimoto H, Morita H,
875 Arad M, Seidman CE, Seidman JG, Ingwall JS, Balschi JA, Tian R. Aberrant activation of
876 AMP-activated protein kinase remodels metabolic network in favor of cardiac glycogen
877 storage. *J. Clin. Invest.* 2007;117:1432-1439
- 878 83. Kim M, Hunter RW, Garcia-Menendez L, Gong G, Yang YY, Kolwicz SC, Jr., Xu J, Sakamoto
879 K, Wang W, Tian R. Mutation in the γ 2-subunit of AMP-activated protein kinase
880 stimulates cardiomyocyte proliferation and hypertrophy independent of glycogen
881 storage. *Circ. Res.* 2014;114:966-975
- 882 84. Dalle Pezze P, Ruf S, Sonntag AG, Langelaar-Makkinje M, Hall P, Heberle AM, Razquin
883 Navas P, van Eunen K, Tolle RC, Schwarz JJ, Wiese H, Warscheid B, Deitersen J, Stork B,
884 Fassler E, Schauble S, Hahn U, Horvatovich P, Shanley DP, Thedieck K. A systems study
885 reveals concurrent activation of AMPK and mTOR by amino acids. *Nat. Commun.*
886 2016;7:13254
- 887 85. Arad M, Moskowitz IP, Patel VV, Ahmad F, Perez-Atayde AR, Sawyer DB, Walter M, Li GH,
888 Burgon PG, Maguire CT, Stapleton D, Schmitt JP, Guo XX, Pizard A, Kupersmidt S, Roden
889 DM, Berul CI, Seidman CE, Seidman JG. Transgenic mice overexpressing mutant PRKAG2
890 define the cause of Wolff-Parkinson-White syndrome in glycogen storage
891 cardiomyopathy. *Circulation*. 2003;107:2850-2856
- 892 86. Yavari A, Stocker CJ, Ghaffari S, Wargent ET, Steeples V, Czibik G, Pinter K, Bellahcene M,
893 Woods A, Martinez de Morentin PB, Cansell C, Lam BY, Chuster A, Petkevicius K, Nguyen-
894 Tu MS, Martinez-Sanchez A, Pullen TJ, Oliver PL, Stockenhuber A, Nguyen C, Lazdam M,
895 O'Dowd JF, Harikumar P, Toth M, Beall C, Kyriakou T, Parnis J, Sarma D, Katritsis G,
896 Wortmann DD, Harper AR, Brown LA, Willows R, Gandra S, Poncio V, de Oliveira
897 Figueiredo MJ, Qi NR, Peirson SN, McCrimmon RJ, Gereben B, Tretter L, Fekete C,
898 Redwood C, Yeo GS, Heisler LK, Rutter GA, Smith MA, Withers DJ, Carling D, Sternick EB,
899 Arch JR, Cawthorne MA, Watkins H, Ashrafian H. Chronic activation of γ 2 AMPK
900 induces obesity and reduces beta cell function. *Cell Metab.* 2016;23:821-836
- 901 87. Yang X, Mudgett J, Bou-About G, Champy MF, Jacobs H, Monassier L, Pavlovic G, Sorg T,
902 Herault Y, Petit-Demouliere B, Lu K, Feng W, Wang H, Ma LJ, Askew R, Erion MD, Kelley
903 DE, Myers RW, Li C, Guan HP. Physiological expression of AMPK γ 2RG mutation
904 causes Wolff-Parkinson-White syndrome and induces kidney injury in mice. *J. Biol.*
905 *Chem.* 2016;291:23428-23439

- 906 88. Yang Y, Atasoy D, Su HH, Sternson SM. Hunger states switch a flip-flop memory circuit
907 via a synaptic AMPK-dependent positive feedback loop. *Cell*. 2011;146:992-1003
- 908 89. Andersson U, Filipsson K, Abbott CR, Woods A, Smith K, Bloom SR, Carling D, Small CJ.
909 AMP-activated protein kinase plays a role in the control of food intake. *J. Biol. Chem*.
910 2004;279:12005-12008
- 911 90. Milan D, Jeon JT, Looft C, Amarger V, Robic A, Thelander M, Rogel-Gaillard C, Paul S,
912 Iannuccelli N, Rask L, Ronne H, Lundstrom K, Reinsch N, Gellin J, Kalm E, Roy PL,
913 Chardon P, Andersson L. A mutation in PRKAG3 associated with excess glycogen content
914 in pig skeletal muscle. *Science*. 2000;288:1248-1251
- 915 91. Costford SR, Kavaslar N, Ahituv N, Chaudhry SN, Schackwitz WS, Dent R, Pennacchio LA,
916 McPherson R, Harper ME. Gain-of-function R225W mutation in human AMPKgamma3
917 causing increased glycogen and decreased triglyceride in skeletal muscle. *PLoS ONE*.
918 2007;2:e903
- 919 92. Crawford SA, Costford SR, Aguer C, Thomas SC, deKemp RA, DaSilva JN, Lafontaine D,
920 Kendall M, Dent R, Beanlands RS, McPherson R, Harper ME. Naturally occurring R225W
921 mutation of the gene encoding AMP-activated protein kinase (AMPK)gamma(3) results
922 in increased oxidative capacity and glucose uptake in human primary myotubes.
923 *Diabetologia*. 2010;53:1986-1997
- 924 93. Morrow VA, Fougelle F, Connell JM, Petrie JR, Gould GW, Salt IP. Direct activation of AMP-
925 activated protein kinase stimulates nitric-oxide synthesis in human aortic endothelial
926 cells. *J. Biol. Chem*. 2003;278:31629-31639
- 927 94. Stahmann N, Woods A, Spengler K, Heslegrave A, Bauer R, Krause S, Viollet B, Carling D,
928 Heller R. Activation of AMP-activated protein kinase by vascular endothelial growth
929 factor mediates endothelial angiogenesis independently of nitric-oxide synthase. *J. Biol.*
930 *Chem*. 2010;285:10638-10652
- 931 95. Dong Y, Zhang M, Liang B, Xie Z, Zhao Z, Asfa S, Choi HC, Zou MH. Reduction of AMP-
932 activated protein kinase alpha2 increases endoplasmic reticulum stress and
933 atherosclerosis in vivo. *Circulation*. 2010;121:792-803
- 934 96. Bess E, Fisslthaler B, Fromel T, Fleming I. Nitric oxide-induced activation of the AMP-
935 activated protein kinase alpha2 subunit attenuates IkappaB kinase activity and
936 inflammatory responses in endothelial cells. *PLoS One*. 2011;6:e20848
- 937 97. Wang S, Zhang M, Liang B, Xu J, Xie Z, Liu C, Viollet B, Yan D, Zou MH. AMPK{alpha}2
938 deletion causes aberrant expression and activation of NAD(P)H oxidase and consequent
939 endothelial dysfunction in vivo. Role of 26S proteasomes. *Circ. Res*. 2010
- 940 98. Nagata D, Mogi M, Walsh K. AMP-activated protein kinase (AMPK) signaling in
941 endothelial cells is essential for angiogenesis in response to hypoxic stress. *J. Biol. Chem*.
942 2003;278:31000-31006

99. Wang J, Alexanian A, Ying R, Kizhakekuttu TJ, Dharmashankar K, Vasquez-Vivar J, Gutterman DD, Widlansky ME. Acute exposure to low glucose rapidly induces endothelial dysfunction and mitochondrial oxidative stress: role for AMP kinase. *Arteriosclerosis Thromb. Vasc. Biol.* 2012;32:712-720
100. Fleming I, Fisslthaler B, Dixit M, Busse R. Role of PECAM-1 in the shear-stress-induced activation of Akt and the endothelial nitric oxide synthase (eNOS) in endothelial cells. *J. Cell Sci.* 2005;118:4103-4111
101. Ouchi N, Kobayashi H, Kihara S, Kumada M, Sato K, Inoue T, Funahashi T, Walsh K. Adiponectin stimulates angiogenesis by promoting cross-talk between AMP-activated protein kinase and Akt signaling in endothelial cells. *J. Biol. Chem.* 2004;279:1304-1309
102. Day RM, Lee YH, Han L, Kim YC, Feng YH. Angiotensin II activates AMPK for execution of apoptosis through energy-dependent and -independent mechanisms. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 2011;301:L772-L781
103. Xu X, Jhun BS, Ha CH, Jin ZG. Molecular mechanisms of ghrelin-mediated endothelial nitric oxide synthase activation. *Endocrinology.* 2008;149:4183-4192
104. Stahmann N, Woods A, Carling D, Heller R. Thrombin activates AMP-activated protein kinase in endothelial cells via a pathway involving Ca²⁺/calmodulin-dependent protein kinase kinase beta. *Mol. Cell. Biol.* 2006;26:5933-5945
105. Levine YC, Li GK, Michel T. Agonist-modulated regulation of AMP-activated protein kinase (AMPK) in endothelial cells. Evidence for an AMPK -> Rac1 -> Akt -> endothelial nitric-oxide synthase pathway. *J. Biol. Chem.* 2007;282:20351-20364
106. Mount PF, Lane N, Venkatesan S, Steinberg GR, Fraser SA, Kemp BE, Power DA. Bradykinin stimulates endothelial cell fatty acid oxidation by CaMKK-dependent activation of AMPK. *Atherosclerosis.* 2008;200:28-36
107. Yang S, Wang J. Estrogen activates AMP-activated protein kinase in human endothelial cells via ERbeta/Ca/calmodulin-dependent protein kinase kinase beta pathway. *Cell biochemistry and biophysics.* 2015;72:701-707
108. Reihill JA, Ewart MA, Salt IP. The role of AMP-activated protein kinase in the functional effects of vascular endothelial growth factor-A and -B in human aortic endothelial cells. *Vasc. Cell.* 2011;3:9
109. Ceolotto G, Gallo A, Papparella I, Franco L, Murphy E, Iori E, Pagnin E, Fadini GP, Albiero M, Semplicini A, Avogaro A. Rosiglitazone reduces glucose-induced oxidative stress mediated by NAD(P)H oxidase via AMPK-dependent mechanism. *Arterioscler. Thromb. Vasc. Biol.* 2007;27:2627-2633
110. Rutherford C, Speirs C, Williams JJ, Ewart MA, Mancini SJ, Hawley SA, Delles C, Viollet B, Costa-Pereira AP, Baillie GS, Salt IP, Palmer TM. Phosphorylation of Janus kinase 1 (JAK1) by AMP-activated protein kinase (AMPK) links energy sensing to anti-inflammatory signaling. *Sci. Signal.* 2016;9:ra109

111. Hwang HJ, Chung HS, Jung TW, Ryu JY, Hong HC, Seo JA, Kim SG, Kim NH, Choi KM, Choi DS, Baik SH, Yoo HJ. The dipeptidyl peptidase-IV inhibitor inhibits the expression of vascular adhesion molecules and inflammatory cytokines in HUVECs via Akt- and AMPK-dependent mechanisms. *Mol. Cell. Endocrinol.* 2015;405:25-34
112. Krasner NM, Ido Y, Ruderman NB, Cacicedo JM. Glucagon-like peptide-1 (GLP-1) analog liraglutide inhibits endothelial cell inflammation through a calcium and AMPK dependent mechanism. *PLoS One.* 2014;9:e97554
113. Sun W, Lee TS, Zhu M, Gu C, Wang Y, Zhu Y, Shyy JY. Statins activate AMP-activated protein kinase in vitro and in vivo. *Circulation.* 2006;114:2655-2662
114. Murakami H, Murakami R, Kambe F, Cao X, Takahashi R, Asai T, Hirai T, Numaguchi Y, Okumura K, Seo H, Murohara T. Fenofibrate activates AMPK and increases eNOS phosphorylation in HUVEC. *Biochem. Biophys. Res. Commun.* 2006;341:973-978
115. Wu Y, Song P, Xu J, Zhang M, Zou MH. Activation of protein phosphatase 2A by palmitate inhibits AMP-activated protein kinase. *J. Biol. Chem.* 2007;282:9777-9788
116. Weikel KA, Cacicedo JM, Ruderman NB, Ido Y. Glucose and palmitate uncouple AMPK from autophagy in human aortic endothelial cells. *Am. J. Physiol. Cell Physiol.* 2015;308:C249-C263
117. Cacicedo JM, Yagihashi N, Keaney JF, Jr., Ruderman NB, Ido Y. AMPK inhibits fatty acid-induced increases in NF-kappaB transactivation in cultured human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* 2004;324:1204-1209
118. Heathcote HR, Mancini SJ, Strembitska A, Jamal K, Reihill JA, Palmer TM, Gould GW, Salt IP. Protein kinase C phosphorylates AMP-activated protein kinase alpha1 Ser487. *Biochem. J.* 2016;473:4681-4697
119. Dagher Z, Ruderman N, Tornheim K, Ido Y. The effect of AMP-activated protein kinase and its activator AICAR on the metabolism of human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* 1999;265:112-115
120. Ido Y, Carling D, Ruderman N. Hyperglycemia-induced apoptosis in human umbilical vein endothelial cells: inhibition by the AMP-activated protein kinase activation. *Diabetes.* 2002;51:159-167
121. Goirand F, Solar M, Athea Y, Viollet B, Mateo P, Fortin D, Leclerc J, Hoerter J, Ventura-Clapier R, Garnier A. Activation of AMP kinase alpha1 subunit induces aortic vasorelaxation in mice. *J. Physiol.* 2007;581:1163-1171
122. Song P, Wang S, He C, Liang B, Viollet B, Zou MH. AMPK-alpha2 deletion exacerbates neointima formation by upregulating Skp2 in vascular smooth muscle cells. *Circ. Res.* 2011;109:1230-1239
123. Motobayashi Y, Izawa-Ishizawa Y, Ishizawa K, Orino S, Yamaguchi K, Kawazoe K, Hamano S, Tsuchiya K, Tomita S, Tamaki T. Adiponectin inhibits insulin-like growth factor-1-induced cell migration by the suppression of extracellular signal-regulated kinase 1/2 activation, but not Akt in vascular smooth muscle cells. *Hypertens. Res.* 2009;32:188-193

124. Ning J, Clemmons DR. AMP-activated protein kinase inhibits IGF-I signaling and protein synthesis in vascular smooth muscle cells via stimulation of insulin receptor substrate 1 S794 and tuberous sclerosis 2 S1345 phosphorylation. *Mol. Endocrinol.* 2010;24:1218-1229
125. Gayard M, Guilluy C, Rousselle A, Viollet B, Henrion D, Pacaud P, Loirand G, Rollier-Derkinderen M. AMPK alpha 1-induced RhoA phosphorylation mediates vasoprotective effect of estradiol. *Arterioscler. Thromb. Vasc. Biol.* 2011;31:2634-2642
126. Pyla R, Pichavaram P, Fairaq A, Park MA, Kozak M, Kamath V, Patel VS, Segar L. Altered energy state reversibly controls smooth muscle contractile function in human saphenous vein during acute hypoxia-reoxygenation: Role of glycogen, AMP-activated protein kinase, and insulin-independent glucose uptake. *Biochem. Pharmacol.* 2015;97:77-88
127. Nagata D, Takeda R, Sata M, Satonaka H, Suzuki E, Nagano T, Hirata Y. AMP-activated protein kinase inhibits angiotensin II-stimulated vascular smooth muscle cell proliferation. *Circulation.* 2004;110:444-451
128. Kim JE, Choi HC. Losartan Inhibits vascular smooth muscle cell proliferation through activation of AMP-activated protein kinase. *The Korean journal of physiology & pharmacology : official journal of the Korean Physiological Society and the Korean Society of Pharmacology.* 2010;14:299-304
129. Ning J, Xi G, Clemmons DR. Suppression of AMPK activation via S485 phosphorylation by IGF-I during hyperglycemia is mediated by AKT activation in vascular smooth muscle cells. *Endocrinology.* 2011;152:3143-3154
130. Son BK, Akishita M, Iijima K, Kozaki K, Maemura K, Eto M, Ouchi Y. Adiponectin antagonizes stimulatory effect of tumor necrosis factor-alpha on vascular smooth muscle cell calcification: regulation of growth arrest-specific gene 6-mediated survival pathway by adenosine 5'-monophosphate-activated protein kinase. *Endocrinology.* 2008;149:1646-1653
131. Cao X, Li H, Tao H, Wu N, Yu L, Zhang D, Lu X, Zhu J, Lu Z, Zhu Q. Metformin inhibits vascular calcification in female rat aortic smooth muscle cells via the AMPK-eNOS-NO pathway. *Endocrinology.* 2013;154:3680-3689
132. Siragusa M, Fleming I. The eNOS signalosome and its link to endothelial dysfunction. *Pflugers Arch.* 2016;468:1125-1137
133. Chen ZP, Mitchelhill KI, Michell BJ, Stapleton D, Rodriguez-Crespo I, Witters LA, Power DA, Ortiz de Montellano PR, Kemp BE. AMP-activated protein kinase phosphorylation of endothelial NO synthase. *FEBS Lett.* 1999;443:285-289
134. Chen Z, Peng IC, Sun W, Su MI, Hsu PH, Fu Y, Zhu Y, Defea K, Pan S, Tsai MD, Shyy JY. AMP-activated protein kinase functionally phosphorylates endothelial nitric oxide synthase Ser633. *Circ. Res.* 2009;104:496-505

- 1059 135. Wang S, Xu J, Song P, Viollet B, Zou MH. In vivo activation of AMP-activated protein
1060 kinase attenuates diabetes-enhanced degradation of GTP cyclohydrolase I. *Diabetes*.
1061 2009;58:1893-1901
- 1062 136. Mount PF, Hill RE, Fraser SA, Levidiotis V, Katsis F, Kemp BE, Power DA. Acute renal
1063 ischemia rapidly activates the energy sensor AMPK but does not increase
1064 phosphorylation of eNOS-Ser1177. *Am. J. Physiol. Renal Physiol.* 2005;289:F1103-F1115
- 1065 137. Kukidome D, Nishikawa T, Sonoda K, Imoto K, Fujisawa K, Yano M, Motoshima H,
1066 Taguchi T, Matsumura T, Araki E. Activation of AMP-activated protein kinase reduces
1067 hyperglycemia-induced mitochondrial reactive oxygen species production and promotes
1068 mitochondrial biogenesis in human umbilical vein endothelial cells. *Diabetes*.
1069 2006;55:120-127
- 1070 138. Shah MS, Brownlee M. Molecular and cellular mechanisms of cardiovascular disorders in
1071 diabetes. *Circ. Res.* 2016;118:1808-1829
- 1072 139. Colombo SL, Moncada S. AMPK α 1 regulates the antioxidant status of vascular
1073 endothelial cells. *Biochem. J.* 2009;421:163-169
- 1074 140. Hou X, Song J, Li XN, Zhang L, Wang X, Chen L, Shen YH. Metformin reduces intracellular
1075 reactive oxygen species levels by upregulating expression of the antioxidant thioredoxin
1076 via the AMPK-FOXO3 pathway. *Biochem. Biophys. Res. Commun.* 2010;396:199-205
- 1077 141. Dong Y, Zhang M, Wang S, Liang B, Zhao Z, Liu C, Wu M, Choi HC, Lyons TJ, Zou MH.
1078 Activation of AMP-activated protein kinase inhibits oxidized LDL-triggered endoplasmic
1079 reticulum stress in vivo. *Diabetes*. 2010;59:1386-1396
- 1080 142. Ridker PM, Luscher TF. Anti-inflammatory therapies for cardiovascular disease. *Eur.*
1081 *Heart J.* 2014;35:1782-1791
- 1082 143. Salt IP, Palmer TM. Exploiting the anti-inflammatory effects of AMP-activated protein
1083 kinase activation. *Expert Opin. Investig. Drugs.* 2012;21:1155-1167
- 1084 144. Ewart MA, Kohlhaas CF, Salt IP. Inhibition of tumor necrosis factor α -stimulated
1085 monocyte adhesion to human aortic endothelial cells by AMP-activated protein kinase.
1086 *Arterioscler. Thromb. Vasc. Biol.* 2008;28:2255-2257
- 1087 145. Spiecker M, Peng HB, Liao JK. Inhibition of endothelial vascular cell adhesion molecule-1
1088 expression by nitric oxide involves the induction and nuclear translocation of
1089 I κ B α . *J. Biol. Chem.* 1997;272:30969-30974
- 1090 146. Zhang Y, Qiu J, Wang X, Xia M. AMP-activated protein kinase suppresses endothelial cell
1091 inflammation through phosphorylation of transcriptional coactivator p300. *Arterioscler.*
1092 *Thromb. Vasc. Biol.* 2011;31:2897-2908
- 1093 147. Schulz E, Doppeide J, Schuhmacher S, Thomas SR, Chen K, Daiber A, Wenzel P, Munzel T,
1094 Keaney JF, Jr. Suppression of the JNK pathway by induction of a metabolic stress
1095 response prevents vascular injury and dysfunction. *Circulation.* 2008;118:1347-1357

148. Mancini SJ, White AD, Bijland S, Rutherford C, Graham D, Richter EA, Viollet B, Touyz RM, Palmer TM, Salt IP. Activation of AMP-activated protein kinase rapidly suppresses multiple pro-inflammatory pathways in adipocytes including IL-1 receptor-associated kinase-4 phosphorylation. *Mol. Cell. Endocrinol.* 2017;440:44-56
149. Kim SA, Choi HC. Metformin inhibits inflammatory response via AMPK-PTEN pathway in vascular smooth muscle cells. *Biochem. Biophys. Res. Commun.* 2012;425:866-872
150. He C, Li H, Viollet B, Zou MH, Xie Z. AMPK suppresses vascular inflammation in vivo by inhibiting Signal Transducer and Activator of Transcription-1. *Diabetes.* 2015;64:4285-4297
151. Zippel N, Malik RA, Fromel T, Popp R, Bess E, Strilic B, Wettschureck N, Fleming I, Fisslthaler B. Transforming growth factor-beta-activated kinase 1 regulates angiogenesis via AMP-activated protein kinase-alpha1 and redox balance in endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* 2013;33:2792-2799
152. Xu MJ, Song P, Shirwany N, Liang B, Xing J, Viollet B, Wang X, Zhu Y, Zou MH. Impaired expression of uncoupling protein 2 causes defective postischemic angiogenesis in mice deficient in AMP-activated protein kinase alpha subunits. *Arterioscler. Thromb. Vasc. Biol.* 2011;31:1757-1765
153. Ouchi N, Shibata R, Walsh K. AMP-activated protein kinase signaling stimulates VEGF expression and angiogenesis in skeletal muscle. *Circ. Res.* 2005;96:838-846
154. Wang S, Liang B, Viollet B, Zou MH. Inhibition of the AMP-activated protein kinase-alpha2 accentuates agonist-induced vascular smooth muscle contraction and high blood pressure in mice. *Hypertension.* 2011;57:1010-1017
155. Schneider H, Schubert KM, Blodow S, Kreutz CP, Erdogmus S, Wiedenmann M, Qiu J, Fey T, Ruth P, Lubomirov LT, Pfitzer G, Mederos Y, M. S, Hardie DG, Gudermann T, Pohl U. AMPK dilates resistance arteries via activation of SERCA and BKCa channels in smooth muscle. *Hypertension.* 2015;66:108-116
156. Igata M, Motoshima H, Tsuruzoe K, Kojima K, Matsumura T, Kondo T, Taguchi T, Nakamaru K, Yano M, Kukidome D, Matsumoto K, Toyonaga T, Asano T, Nishikawa T, Araki E. Adenosine monophosphate-activated protein kinase suppresses vascular smooth muscle cell proliferation through the inhibition of cell cycle progression. *Circ. Res.* 2005;97:837-844
157. Song P, Wang S, He C, Liang B, Viollet B, Zou MH. AMPKalpha2 deletion exacerbates neointima formation by upregulating Skp2 in vascular smooth muscle cells. *Circ. Res.* 2011;109:1230-1239
158. Ding Y, Zhang M, Zhang W, Lu Q, Cai Z, Song P, Okon IS, Xiao L, Zou MH. AMP-activated protein kinase alpha 2 deletion induces VSMC phenotypic switching and reduces features of atherosclerotic plaque stability. *Circ. Res.* 2016;119:718-730

159. Song P, Zhou Y, Coughlan KA, Dai X, Xu H, Viollet B, Zou MH. Adenosine monophosphate-activated protein kinase- α 2 deficiency promotes vascular smooth muscle cell migration via S-phase kinase-associated protein 2 upregulation and E-cadherin downregulation. *Arterioscler. Thromb. Vasc. Biol.* 2013;33:2800-2809
160. Cai Z, Ding Y, Zhang M, Lu Q, Wu S, Zhu H, Song P, Zou MH. Ablation of adenosine monophosphate-activated protein kinase α 1 in vascular smooth muscle cells promotes diet-Induced atherosclerotic calcification in vivo. *Circ. Res.* 2016;119:422-433
161. Cacicedo JM, Gauthier MS, Lebrasseur NK, Jasuja R, Ruderman NB, Ido Y. Acute exercise activates AMPK and eNOS in the mouse aorta. *Am. J. Physiol. Heart Circ. Physiol.* 2011;301:H1255-H1265
162. Ma L, Ma S, He H, Yang D, Chen X, Luo Z, Liu D, Zhu Z. Perivascular fat-mediated vascular dysfunction and remodeling through the AMPK/mTOR pathway in high-fat diet-induced obese rats. *Hypertens. Res.*
163. Zhao CX, Xu X, Cui Y, Wang P, Wei X, Yang S, Edin ML, Zeldin DC, Wang DW. Increased endothelial nitric-oxide synthase expression reduces hypertension and hyperinsulinemia in fructose-treated rats. *J. Pharmacol. Exp. Ther.* 2009;328:610-620
164. Bradley EA, Eringa EC, Stehouwer CD, Korstjens I, van Nieuw Amerongen GP, Musters R, Sipkema P, Clark MG, Rattigan S. Activation of AMP-activated protein kinase by 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside in the muscle microcirculation increases nitric oxide synthesis and microvascular perfusion. *Arterioscler. Thromb. Vasc. Biol.* 2010;30:1137-1142
165. Ford RJ, Teschke SR, Reid EB, Durham KK, Kroetsch JT, Rush JW. AMP-activated protein kinase activator AICAR acutely lowers blood pressure and relaxes isolated resistance arteries of hypertensive rats. *J. Hypertens.* 2012;30:725-733
166. Ford RJ, Rush JW. Endothelium-dependent vasorelaxation to the AMPK activator AICAR is enhanced in aorta from hypertensive rats and is NO and EDCF dependent. *Am. J. Physiol. Heart Circ. Physiol.* 2011;300:H64-H75
167. Bosselaar M, Boon H, van Loon LJ, van den Broek PH, Smits P, Tack CJ. Intra-arterial AICA-riboside administration induces NO-dependent vasodilation in vivo in human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 2009;297:E759-E766
168. Enkhjargal B, Godo S, Sawada A, Suvd N, Saito H, Noda K, Satoh K, Shimokawa H. Endothelial AMP-activated protein kinase regulates blood pressure and coronary flow responses through hyperpolarization mechanism in mice. *Arterioscler. Thromb. Vasc. Biol.* 2014;34:1505-1513
169. Liang B, Wang S, Wang Q, Zhang W, Viollet B, Zhu Y, Zou MH. Aberrant endoplasmic reticulum stress in vascular smooth muscle increases vascular contractility and blood pressure in mice deficient of AMP-activated protein kinase- α 2 in vivo. *Arterioscler. Thromb. Vasc. Biol.* 2013;33:595-604

170. Kohlstedt K, Trouvain C, Boettger T, Shi L, Fisslthaler B, Fleming I. AMP-activated protein kinase regulates endothelial cell angiotensin-converting enzyme expression via p53 and the post-transcriptional regulation of microRNA-143/145. *Circ. Res.* 2013;112:1150-1158
171. Buhl ES, Jessen N, Pold R, Ledet T, Flyvbjerg A, Pedersen SB, Pedersen O, Schmitz O, Lund S. Long-term AICAR administration reduces metabolic disturbances and lowers blood pressure in rats displaying features of the insulin resistance syndrome. *Diabetes.* 2002;51:2199-2206
172. Meijer RJ, Bakker W, Alta CL, Sipkema P, Yudkin JS, Viollet B, Richter EA, Smulders YM, van Hinsbergh VW, Serne EH, Eringa EC. Perivascular adipose tissue control of insulin-induced vasoreactivity in muscle is impaired in db/db mice. *Diabetes.* 2013;62:590-598
173. Almabrouk TA, Ugusman AB, Katwan OJ, Salt IP, Kennedy S. Deletion of AMPK α 1 attenuates the anticontractile effect of perivascular adipose tissue (PVAT) and reduces adiponectin release. *Br. J. Pharmacol.* 2017;in press
174. Gaskin FS, Kamada K, Zuidema MY, Jones AW, Rubin LJ, Korthuis RJ. Isoform-selective 5'-AMP-activated protein kinase-dependent preconditioning mechanisms to prevent postischemic leukocyte-endothelial cell adhesive interactions. *Am. J. Physiol. Heart Circ. Physiol.* 2011;300:H1352-H1360
175. Li D, Wang D, Wang Y, Ling W, Feng X, Xia M. Adenosine monophosphate-activated protein kinase induces cholesterol efflux from macrophage-derived foam cells and alleviates atherosclerosis in apolipoprotein E-deficient mice. *J. Biol. Chem.* 2010;285:33499-33509
176. Chen B, Li J, Zhu H. AMP-activated protein kinase attenuates oxLDL uptake in macrophages through PP2A/NF-kappaB/LOX-1 pathway. *Vascul. Pharmacol.* 2016;85:1-10
177. Vasamsetti SB, Karnewar S, Kanugula AK, Thatipalli AR, Kumar JM, Kotamraju S. Metformin inhibits monocyte-to-macrophage differentiation via AMPK-mediated inhibition of STAT3 Activation: potential role in atherosclerosis. *Diabetes.* 2015;64:2028-2041
178. Wang Q, Zhang M, Liang B, Shirwany N, Zhu Y, Zou MH. Activation of AMP-activated protein kinase is required for berberine-induced reduction of atherosclerosis in mice: the role of uncoupling protein 2. *PLoS One.* 2011;6:e25436
179. UKPDS. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). UK Prospective Diabetes Study (UKPDS) Group. *Lancet.* 1998;352:854-865
180. Dai X, Ding Y, Liu Z, Zhang W, Zou MH. Phosphorylation of CHOP (C/EBP Homologous Protein) by the AMP-activated protein kinase α 1 in macrophages promotes CHOP degradation and reduces injury-induced neointimal disruption in vivo. *Circ. Res.* 2016;119:1089-1100

- 1210 181. Li FY, Lam KS, Tse HF, Chen C, Wang Y, Vanhoutte PM, Xu A. Endothelium-selective
1211 activation of AMP-activated protein kinase prevents diabetes mellitus-induced
1212 impairment in vascular function and reendothelialization via induction of heme
1213 oxygenase-1 in mice. *Circulation*. 2012;126:1267-1277
- 1214 182. Cao Q, Cui X, Wu R, Zha L, Wang X, Parks JS, Yu L, Shi H, Xue B. Myeloid deletion of
1215 alpha1AMPK exacerbates atherosclerosis in LDL receptor knockout (LDLRKO) mice.
1216 *Diabetes*. 2016;65:1565-1576
- 1217 183. Song Y, Wu Y, Su X, Zhu Y, Liu L, Pan Y, Zhu B, Yang L, Gao L, Li M. Activation of AMPK
1218 inhibits PDGF-induced pulmonary arterial smooth muscle cells proliferation and its
1219 potential mechanisms. *Pharmacol. Res.* 2016;107:117-124
- 1220 184. Dean A, Nilsen M, Loughlin L, Salt IP, MacLean MR. Metformin reverses development of
1221 pulmonary hypertension via aromatase inhibition. *Hypertension*. 2016;68:446-454
- 1222 185. Omura J, Satoh K, Kikuchi N, Satoh T, Kurosawa R, Nogi M, Otsuki T, Kozu K, Numano K,
1223 Suzuki K, Sunamura S, Tatebe S, Aoki T, Sugimura K, Miyata S, Hoshikawa Y, Okada Y,
1224 Shimokawa H. Protective roles of endothelial AMP-activated protein kinase against
1225 hypoxia-induced pulmonary hypertension in mice. *Circ. Res.* 2016;119:197-209
- 1226 186. Ibe JC, Zhou Q, Chen T, Tang H, Yuan JX, Raj JU, Zhou G. Adenosine monophosphate-
1227 activated protein kinase is required for pulmonary artery smooth muscle cell survival
1228 and the development of hypoxic pulmonary hypertension. *Am. J. Respir. Cell. Mol. Biol.*
1229 2013;49:609-618
- 1230 187. Moral-Sanz J, Mahmoud AD, Ross FA, Eldstrom J, Fedida D, Hardie DG, Evans AM. AMP-
1231 activated protein kinase inhibits Kv 1.5 channel currents of pulmonary arterial myocytes
1232 in response to hypoxia and inhibition of mitochondrial oxidative phosphorylation. *J.*
1233 *Physiol.* 2016;594:4901-4915
- 1234 188. Pinter K, Grignani RT, Czibik G, Farza H, Watkins H, Redwood C. Embryonic expression
1235 of AMPK gamma subunits and the identification of a novel gamma2 transcript variant in
1236 adult heart. *J. Mol. Cell. Cardiol.* 2012;53:342-349
- 1237 189. Cool B, Zinker B, Chiou W, Kifle L, Cao N, Perham M, Dickinson R, Adler A, Gagne G,
1238 Iyengar R, Zhao G, Marsh K, Kym P, Jung P, Camp HS, Frevert E. Identification and
1239 characterization of a small molecule AMPK activator that treats key components of type
1240 2 diabetes and the metabolic syndrome. *Cell Metab.* 2006;3:403-416
- 1241 190. Zadra G, Photopoulos C, Tyekucheva S, Heidari P, Weng QP, Fedele G, Liu H, Scaglia N,
1242 Priolo C, Sicinska E, Mahmood U, Signoretti S, Birnberg N, Loda M. A novel direct
1243 activator of AMPK inhibits prostate cancer growth by blocking lipogenesis. *EMBO Mol.*
1244 *Med.* 2014;6:519-538

- 1245 191. Salatto CT, Miller RA, Cameron KO, Cokorinos E, Reyes A, Ward J, Calabrese M,
1246 Kurumbail R, Rajamohan F, Kalgutkar AS, Tess DA, Shavnya A, Genung NE, Edmonds DJ,
1247 Jatkar A, Maciejewski BS, Amaro M, Gandhok H, Monetti M, Cialdea K, Bollinger E,
1248 Kreeger JM, Coskran TM, Opsahl AC, Boucher GG, Birnbaum MJ, DaSilva-Jardine P, Rolph
1249 T. Selective Activation of AMPK b1-containing Isoforms Improves Kidney Function in a
1250 Rat Model of Diabetic Nephropathy. *J. Pharmacol. Exp. Ther.* 2017
- 1251 192. Gollob MH, Green MS, Tang AS, Gollob T, Karibe A, Ali Hassan AS, Ahmad F, Lozado R,
1252 Shah G, Fananapazir L, Bachinski LL, Roberts R. Identification of a gene responsible for
1253 familial Wolff-Parkinson-White syndrome. *N. Engl. J. Med.* 2001;344:1823-1831
- 1254 193. Arad M, D.W. B, Perez-Atayde AR, McKenna WJ, Sparks EA, Kanter RJ, McGarry K,
1255 Seidman JG, Seidman CE. Constitutively active AMP kinase mutations cause glycogen
1256 storage disease mimicking hypertrophic cardiomyopathy. *J. Clin. Invest.* 2002;109:357-
1257 362
- 1258 194. Charron P, Genest M, Richard P, Komajda M, Pochmalicki G. A familial form of
1259 conduction defect related to a mutation in the PRKAG2 gene. *Europace.* 2007;9:597-600
- 1260 195. Blair E, Redwood C, Ashrafian H, Oliveira M, Broxholme J, Kerr B, Salmon A, Ostman-
1261 Smith I, Watkins H. Mutations in the gamma-2 subunit of AMP-activated protein kinase
1262 cause familial hypertrophic cardiomyopathy: evidence for the central role of energy
1263 compromise in disease pathogenesis. *Hum. Mol. Genet.* 2001;10:1215-1220
- 1264 196. Arad M, Maron BJ, Gorham JM, Johnson WH, Jr., Saul JP, Perez-Atayde AR, Spirito P,
1265 Wright GB, Kanter RJ, Seidman CE, Seidman JG. Glycogen storage diseases presenting as
1266 hypertrophic cardiomyopathy. *N. Engl. J. Med.* 2005;352:362-372
- 1267 197. Bayrak F, Komurcu-Bayrak E, Mutlu B, Kahveci G, Basaran Y, Erginel-Unaltuna N.
1268 Ventricular pre-excitation and cardiac hypertrophy mimicking hypertrophic
1269 cardiomyopathy in a Turkish family with a novel PRKAG2 mutation. *Eur. J. Heart Fail.*
1270 2006;8:712-715
- 1271 198. Xie C, Zhang YP, Song L, Luo J, Qi W, Hu J, Lu D, Yang Z, Zhang J, Xiao J, Zhou B, Du JL, Jing
1272 N, Liu Y, Wang Y, Li BL, Song BL, Yan Y. Genome editing with CRISPR/Cas9 in postnatal
1273 mice corrects PRKAG2 cardiac syndrome. *Cell Res.* 2016;26:1099-1111
- 1274 199. Laforet P, Richard P, Said MA, Romero NB, Lacene E, Leroy JP, Baussan C, Hogrel JY,
1275 Lavergne T, Wahbi K, Hainque B, Duboc D. A new mutation in PRKAG2 gene causing
1276 hypertrophic cardiomyopathy with conduction system disease and muscular
1277 glycogenosis. *Neuromuscul. Disord.* 2006;16:178-182
- 1278

FIGURE LEGENDS

Figure 1: Domain layouts of AMPK subunits and their isoforms. The linear layout of domains is shown, approximately to scale and with similar color coding as in Fig. 2. Note that both β subunits are N-myristoylated, and that the $\gamma 2$ and $\gamma 3$ subunit isoforms have unrelated N-terminal extensions of unknown function, although both are also reported to exist as shorter, N-terminally truncated versions due to alternate start sites and/or splicing¹⁸⁸.

Figure 2: Structure of human AMPK ($\alpha 1\beta 2\gamma 1$ complex)⁷. The model was created in spacefilling mode using PyMol v1.7.4.2 with the co-ordinates in PDB file 4RER, and with color coding similar to Fig. 1. The heterotrimer was crystallized in the presence of β -cyclodextrin, which occupies the glycogen-binding site, staurosporine, which occupies the active site, and AMP, which occupies sites 1, 3 and 4 on the γ subunit (sites 1 and 4 are round the back in this view). Although Thr172 was phosphorylated, it is not visible in this view but lies in the cleft between the α subunit C lobe and the β -CTD, just over the right-hand “shoulder” of the C-lobe.

Figure 3: Two views of the structure of the four CBS repeats of the $\gamma 1$ subunit. The model used the co-ordinates in PDB file 2V8Q¹⁸, and was rendered in PyMol v1.7.4.2 with the $\gamma 1$ subunit in cartoon view. The two views are rotated 180° around the x axis (dashed line) with respect to each other, with the orientation of the top view being similar to that in Fig. 2. Note the pseudosymmetrical layout of the four CBS repeats, which are colored differently (and differently to Figs. 1 and 2). Residues equivalent to those mutated in $\gamma 2$ are highlighted using the “dots” version of space-filling representation, and are numbered using the human $\gamma 2$ numbering. The three molecules of bound AMP are labelled and are shown in standard space-filling view, with C atoms green, O red and N blue.

Figure 4: Regulation of endothelial NO and superoxide synthesis by AMPK. AMPK activation stimulates NO synthesis via multiple mechanisms: (i) phosphorylation of eNOS at Ser633 and Ser1177; (ii) increasing Hsp90 association with eNOS; (iii) increasing BH₄ concentrations via GTP cyclohydrolase I (GTPCH1); (iv) reducing superoxide synthesis via inhibition of Nox and increasing antioxidant protein

(superoxide dismutase (SOD), thioredoxin and catalase) levels. NO itself is also reported to activate AMPK.

Figure 5: Actions of AMPK in vascular cells. Physiological activators of AMPK in ECs include hypoxia/ischemia, shear stress, adiponectin, thrombin, bradykinin and VEGF (synthesis of which is stimulated by AMPK in other tissues). AMPK is reported to stimulate VSMC relaxation through: (i) increased NO synthesis and possibly endothelium-dependent hypopolarising factor (EDHF); (ii) inhibition of MYPT1/MLC phosphorylation and Ca^{2+} levels in VSMCs, reported to be mediated by reduced RhoA activity and increased sarco/endoplasmic Ca^{2+} ATPase (SERCA) activity respectively. AMPK activation in ECs stimulates proliferation and migration, whereas in VSMCs, proliferation and migration are inhibited, associated with p53 phosphorylation, Rb dephosphorylation and p27(Kip1) stabilisation. AMPK also inhibits VSMC calcification by reducing Runx2 and pro-inflammatory signalling pathways leading to leukocyte adhesion and cytokine/chemokine synthesis.

Table 1: List of pharmacological agents commonly used to activate AMPK in intact cells or in vivo, and their mechanisms of action.

Class	Agent	Mechanism	Binding site used	Isoform-selective?	Ref.
1	metformin	mitochondrial inhibitor, AMP ↑	γ subunit (binds AMP)	No	33
1	phenformin	mitochondrial inhibitor, AMP ↑	γ subunit (binds AMP)	No	33
1	berberine	mitochondrial inhibitor, AMP ↑	γ subunit (binds AMP)	No	33
2	2-deoxyglucose	glycolytic inhibitor, AMP ↑	γ subunit (binds AMP)	No	33
3	AICAR	pro-drug, converted to ZMP	γ subunit (binds ZMP)	No	33, 34
4	C13	pro-drug, converted to C2	γ subunit (binds C2)	α1-selective	38, 40
5	A769662	direct activator	ADaM site	β1-selective	189
5	991	direct activator	ADaM site	β1>β2	8
5	MT 63-78	direct activator	ADaM site	β1-selective	190
5	PF-06409577	direct activator	ADaM site	β1-selective	191
5	PF-249	direct activator	ADaM site	β1-selective	191
5	salicylate	direct activator	ADaM site	β1-selective	9, 44

Table 2: Amino acid replacements, generated by mutations in *PRKAG2*, that are associated with heart disorders. Equivalent residues in the sequences of human $\gamma 1$ and $\gamma 3$ are shown in the second and third column and the CBS repeat affected in the fourth. V336 and R350 are located in the linker between CBS1 and CBS2.

$\gamma 2$ mutation	$\gamma 1$ equivalent	$\gamma 3$ equivalent	CBS repeat affected	Reference
R302Q	R70	R225	CBS1	192-194
S333P	S101	S333	CBS1	79
V336A	V104	V259	linker	79
L insert (after R350)	after R118	after R273	linker	195
H383R	H151	H306	CBS2	195
R384T	R152	R307	CBS2	77
T400N	T168	T323	CBS2	193
Y487H	Y255	Y410	CBS3	196
N488I	N256	N411	CBS3	193
E506K	E274	E429	CBS4	79, 197
H530R	H298	H453	CBS4	198
R531G	R299	R454	CBS4	80
R531Q	R299	R454	CBS4	78
S548P	S316	S471	CBS4	199









